

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
**WO 01/36979 A2**

- (51) International Patent Classification<sup>7</sup>: **G01N 33/68**
- (21) International Application Number: **PCT/GB00/04315**
- (22) International Filing Date:  
10 November 2000 (10.11.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9927125.6 16 November 1999 (16.11.1999) GB
- (71) Applicant (*for all designated States except US*): **UNIVERSITY OF READING [GB/GB]**; Whiteknights House, Whiteknights, Reading RG6 6AH (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **PAGE, Nigel** [GB/GB]; University of Reading, School of Animal Microbial Sciences, Whiteknights, P.O. Box 228, Reading RG6 6AJ (GB). **LOWRY, Phillip** [GB/GB]; University of Reading, School of Animal and Microbial Sciences, Whiteknights, P.O. Box 228, Reading RG6 6AJ (GB).
- (74) Agents: **CORNISH, K., V., J. et al.**; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— *Without international search report and to be republished upon receipt of that report.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **PLACENTAL HUMAN NEUROKININ B PRECURSOR**

(57) Abstract: Methods of diagnosing pregnancy induced hypertension or pre-eclampsia by the measurement of the production of neurokinin B, its precursor and fragments thereof are provided, as are kits for use in the methods. Treatments of the conditions and methods of preparing suitable medicaments are also provided as are antibodies and useful antigenic materials.

WO 01/36979 A2

## PLACENTAL HUMAN NEUROKININ B PRECURSOR

The present invention is concerned with the detection of the production of the human precursor of neurokinin B by the placenta and to the detection of the production of neurokinin B gene products, or variants, or fragments thereof as a means of predicting the onset of pregnancy induced hypertension or pre-eclampsia or related foetal complications (or following their course). The application is also directed to methods of preventing or treating pregnancy-induced hypertension or pre-eclampsia by suppressing the effects of excessive neurokinin B secreted into maternal blood.

Pregnancy-induced hypertension (PIH) and pre-eclampsia, two of the most elusive and complex conditions of pregnancy, have been very difficult to define and manage. Pre-eclampsia is still one of the most common and life threatening complications of pregnancy in the Western World. The primary cause of pre-eclampsia has been difficult to elucidate because its signs and symptoms have always presented as a cluster of conditions. Hence, it has been defined as a syndrome, commonly presenting with the features of maternal hypertension and proteinuria, but including extensive complications involving the maternal liver, coagulation and nervous systems (Henriksen, T., (1998) Scand. J. Rheumatol. Suppl. 107 86-91). The clinical problems of pre-eclampsia normally become apparent only in the second half of pregnancy and are believed to emerge during the first trimester. It would appear that pre-eclamptic complications only present if placental tissue is present in the uterus of the mother. Indeed, cases of hydatidiform mole can present with pre-eclampsia where the uterus only contains disordered placental tissue (Nugent, C.E, *et al* (1966) Obstet. Gynecol. 87 829-31). Once pre-eclampsia is diagnosed during the course of pregnancy and the placental tissue is surgically removed or expelled during birth the condition ultimately clears. There have been many suggestions about the causes of pre-eclampsia ranging from the development of a poor placental/uterine vascular system to the immunology of incompatibility between the mother

and foetus. Though these theories do have some substance they do not account for the systemic effects of this syndrome. Many symptoms are likely to be the result of secondary effects of hypertension and not the direct cause of the syndrome. Early detection of the development of PIH or pre-eclampsia would therefore be of great benefit in allowing precautionary measures to be taken, including specific treatment of hypertension and other complications associated with pre-eclampsia such as seizures, blot clotting problems etc.

The placental damage visible and hypertension observed in an expectant mother with pre-eclampsia has been implicated in an increased risk of foetal complications including growth retardation and foetal hypoxia. In extreme cases this could be a cause of miscarriage. In other studies, pre-eclampsia has been postulated as a maternal and foetal adaptation to foetal growth retardation. Since not all women with foetal growth retardation develop pre-eclampsia the decisive factor is a maternal response (Walker, J. (2000) The Lancet 356 1260-1265). Characteristics of this adaptation are present in not only pre-eclampsia but also in foetal growth retardation and miscarriage. For example, the failure of the normal expansion of plasma volume in the mother is associated with both impaired foetal growth and pre-eclampsia (Gulmezoglu AM, Hofmeyr GJ (2000) Cochrane Database Syst Rev 2 CD000167). Problems observed in pre-eclampsia such as thrombophilia are suggested to be the result of thrombotic lesions in a pathological placenta (Mousa HA, Alfirevic Z (2000) Hum Reprod 15:1830-3). It is apparent therefore that pre-eclampsia and foetal growth retardation and foetal hypoxia are linked, and diagnostic methods and treatments for pre-eclampsia may also be suitable in the prediction, diagnosis and/or treatment of these foetal conditions.

Neurokinin B (NKB) belongs to a family of peptides called tachykinins, the first and most well known of which is substance P which was discovered in 1931 (von Euler, U.S. and Gaddum, J.H. (1931) J Physiol 72:74-87). It took over another five decades before the discovery of a further two members of

the tachykinin family, one designated substance K or neurokinin A (Kimura, S., *et al* (1983) Proc. Japan Acad 59B 101-104) and the other designated neuromedin K, now know as neurokinin B (Kangawa, K., *et al* (1983). Biochem. Biophys. Res. Commun. 114 533-540). The tachykinins have been

5 implicated to have a wide variety of biological actions from smooth muscle contraction, vasodilation, pain transmission, neurogenic inflammation, to the activation of the immune system (Longmore, J., *et al* (1997) Canadian J. Physio. & Pharmacol. 75 612-621). Neurokinin B has been found to be the

10 most potent neurokinin to cause vasoconstriction of both the mesenteric vascular bed (D'Orleans-Juste, P. *et al* (1991). Eur. J. Pharmacol. 204 329-334) and contraction of the hepatic portal vein (Mastrangelo, D., *et al* (1987) Eur J Pharmacol. 134, 321-6). Neurokinin B is also the most potent member of the family to act at the NK<sub>3</sub> receptor and, whilst substance P and K slow down the heart rate, NK<sub>3</sub> receptor agonists have the opposite effect in that

15 they increase heart rate when perfused in the canine coronary arterial blood supply (Thompson, G.W. *et al* (1998) American Journal of Physiology-Regulatory Integrative and Comparative Physiology 275 (5), 1683-1689). In an animal model, intravenous injections of neurokinin B in guinea pigs have been shown to produce a dose related hypertension, and very high levels of

20 neurokinin B agonist led to animal discomfort (Roccon, A., *et al* (1996) Brit. J. Pharmacol. 118 1095-1102). Similar experiments have shown an increase in blood pressure upon intravenous infusion of neurokinin B in rats (Page *et al.*, (2000) Nature 405 797-800). Neurokinin B has not been reliably found in any peripheral tissues taken from experimental animals; for example, Moussaoui

25 *et al* (Neuroscience (1992) 48, 967-978) tested a wide range of peripheral tissues using a very sensitive and specific assay system and found no trace of neurokinin B at all.

A human neurokinin B precursor has been identified which, on processing,

30 gives rise to a peptide identical to neurokinin B of other mammalian species (bovine, porcine, rat and mouse) (Incyte Pharmaceuticals Inc., International patent application no. WO98/57986). We have discovered, most surprisingly,

that this human neurokinin B precursor is produced by placental tissue during pregnancy and that neurokinin B and fragments of the precursor are passed into the maternal bloodstream.

- 5 We have found that in normal pregnancy, substantial levels (eg 100 picomolar range) of neurokinin B (and other breakdown products of the human neurokinin B precursor) are found in the maternal blood stream near to term, but that zero or very low levels are found before this. However, in some cases near term levels are identified at an early stage of pregnancy (eg  
10 after only 9 weeks), and in cases of pregnancy induced hypertension or pre-eclampsia very high (nanomolar) concentrations of neurokinin B are found in the maternal plasma near to term. Thus, detection of raised plasma levels of neurokinin B, neurokinin B precursor, its breakdown products, or variants thereof at an early stage will provide an indication of the likely development  
15 of pregnancy induced hypertension or pre-eclampsia and may even provide an indication of the likely future severity of these conditions. Furthermore, reduction in the levels of circulating neurokinin B (or reduction of its effects) will ameliorate the adverse effects upon the mother seen in these conditions. As a result of the relationship between pre-eclampsia and foetal  
20 complications including foetal growth retardation and/or foetal hypoxia, neurokinin B agonists or antagonists may be useful in ameliorating these conditions. Overproduction of the human neurokinin B precursor may also be a causative factor in certain hypertensive conditions in non-pregnant individuals (either through the effect of neurokinin B or one or more of the  
25 other breakdown products of the precursor).

- In a first aspect of the invention there is provided a method of predicting pregnancy induced hypertension in a human subject by assessing the concentration in a biological sample, e.g. blood, of a human neurokinin B  
30 precursor gene product or a variant or a fragment thereof.

In a second aspect of the invention there is provided a method of predicting pre-eclampsia or related foetal complications in a human subject by assessing the concentration in a biological sample, e.g. blood, of a human neurokinin B precursor gene product or a variant or a fragment thereof.

5

In a third aspect of the invention there is provided a method of diagnosing pregnancy induced hypertension in a human subject by assessing the concentration in a biological sample, e.g. blood, of a human neurokinin B precursor gene product or a variant or a fragment thereof.

10

In a fourth aspect of the invention there is provided a method of diagnosing pre-eclampsia or related foetal complications in a human subject by assessing the concentration in a biological sample, e.g. blood, of a human neurokinin B precursor gene product or a variant or a fragment thereof.

15

Preferably, the methods of the first, second, third or fourth aspects comprise assessing the concentration in a biological sample, e.g. blood, of neurokinin B.

20 In a fifth aspect of the invention there is provided a method of estimating the likely future degree of pregnancy induced hypertension in a human subject by assessing the concentration in a biological sample, e.g. blood, of human neurokinin B precursor gene product or a variant or a fragment thereof, and correlating the result with the predicted future severity of pregnancy induced  
25 hypertension.

In a sixth aspect of the invention there is provided a method of estimating the likely future degree of pre-eclampsia or related foetal complications in a human subject by assessing the concentration in a biological sample, e.g.  
30 blood, of human neurokinin B precursor or a variant or a fragment thereof, and correlating the result with the predicted future severity of pre-eclampsia or related foetal complications.

Preferably, the methods of the fifth and sixth aspects comprise assessing the concentration in a biological sample, e.g. blood, of neurokinin B, and correlating the result with the predicted future severity of pregnancy induced hypertension or pre-eclampsia or related foetal complications, respectively.

In a seventh aspect of the invention there is provided a method of preventing or treating pregnancy induced hypertension in a human subject by the administration of an agent which inhibits the biological effect of neurokinin B.

In an eighth aspect of the invention there is provided a method of preventing or treating pre-eclampsia or related foetal complications in a human subject by the administration of an agent which inhibits the biological effect of neurokinin B.

In a ninth aspect of the invention there is provided the use of a human neurokinin B precursor gene product or a variant or a fragment thereof in the manufacture of a diagnostic for use in the prediction or diagnosis of pregnancy-induced hypertension.

In a tenth aspect of the invention there is provided the use of a human neurokinin B precursor gene product or a variant or a fragment thereof in the manufacture of a diagnostic for use in the prediction or diagnosis of pre-eclampsia or related foetal complications.

Preferably, the ninth and tenth aspects comprise the use of an epitopic variant or epitopic fragment of human neurokinin B precursor. More preferably, the methods comprise the use of neurokinin B in the manufacture of a diagnostic for use in the prediction or diagnosis of pregnancy induced hypertension, pre-eclampsia or related foetal complications.



In an eleventh aspect of the invention there is provided the use of an agent which inhibits the biological effect of neurokinin B in the manufacture of a medicament for the prevention or treatment of pregnancy induced hypertension.

5

In a twelfth aspect of the invention there is provided the use of an agent which inhibits the biological effect of neurokinin B in the manufacture of a medicament for the prevention or treatment of pre-eclampsia or related foetal complications.

10

In a preferred embodiment of the eleventh and twelfth aspects, there is provided a pharmaceutical composition comprising an agent which inhibits the biological effect of neurokinin B, for use in the prevention or treatment of pregnancy induced hypertension, pre-eclampsia or related foetal complications.

15

In a thirteenth aspect of the invention there is provided a kit for the prediction or diagnosis of pregnancy induced hypertension comprising a binding partner, eg an antibody, to a neurokinin B precursor gene product or variant or fragment thereof.

20

In a fourteenth aspect of the invention there is provided a kit for the prediction or diagnosis of pre-eclampsia or related foetal complications comprising a binding partner, eg an antibody, to a neurokinin B precursor gene product or variant or fragment thereof.

25

In a fifteenth aspect of the invention there is provided a kit for the prediction or diagnosis of pregnancy induced hypertension, comprising a binding partner, eg an antibody, to a neurokinin B precursor gene product or variant or fragment thereof, together with instructions for the performance of an assay for predicting the levels of neurokinin B in a biological sample and

30

correlating the assay results with the likely future development of pregnancy induced hypertension.

In a sixteenth aspect of the invention there is provided a kit for the prediction  
5 or diagnosis of pre-eclampsia or related foetal complications, comprising a  
binding partner, eg an antibody, to neurokinin B precursor gene product or  
variant or fragment thereof, together with instructions for the performance of  
an assay for predicting the levels of neurokinin B in a biological sample and  
correlating the assay results with the likely future development of pre-  
10 eclampsia or related foetal complications.

In a seventeenth aspect of the invention there is provided a kit for use in  
estimating the likely future degree of pregnancy induced hypertension,  
comprising a binding partner, eg an antibody, to a neurokinin B precursor  
15 gene product or variant or fragment thereof, together with instructions for the  
performance of an assay for predicting the levels of neurokinin B in a  
biological sample and correlating the assay results with the predicted future  
severity of pregnancy induced hypertension.

20 In an eighteenth aspect of the invention there is provided a kit for use in  
estimating the likely future degree of pre-eclampsia or related foetal  
complications, comprising a binding partner, eg an antibody, to a neurokinin  
B precursor gene product or variant or fragment thereof, together with  
instructions for the performance of an assay for predicting the levels of  
25 neurokinin B in a biological sample and correlating the assay results with the  
predicted future severity of pre-eclampsia or related foetal complications.

Preferably, the kits of the thirteenth to eighteenth aspects of the invention  
comprise a binding partner, e.g. an antibody, to a neurokinin B precursor,  
30 neurokinin B or epitopic variants or epitopic fragments thereof. More  
preferably the kits comprise a binding partner to the polypeptide sequences  
of Figures 1 or 2, or epitopic variants or epitopic fragments thereof.

In a nineteenth aspect of the invention there is provided the use of an agonist of neurokinin B or neurokinin B in the preparation of a medicament for the reduction of blood volume in cases of hypotension.

5

In a twentieth aspect of the invention there is provided the use of an agonist of neurokinin B or neurokinin B in the reduction of blood volume in cases of hypotension.

- 10 In a twenty-first aspect of the invention there is provided a method of alleviating pre-eclampsia in a human subject by modifying the diet of the human subject to reduce the content of toxin generating substances therein.

- 15 In a twenty-second aspect of the invention there is provided a method of alleviating pre-eclampsia in a human subject including modifying the dietary pattern of the subject to reduce concentrations of potential toxins in the portal vein.

- 20 In a twenty-third aspect of the invention there is provided a dietary methodology for the alleviation of pre-eclampsia in a human subject in which the amount of toxin generating substances is reduced.

Figure 1 shows the polypeptide sequence of cloned human neurokinin B precursor, available under Accession No. aaf76980.

25

Figure 2 shows the polypeptide sequence of the active neurokinin B peptide.

- 30 Figure 3 shows the polynucleotide sequence of placental cDNA of the human neurokinin B precursor, where ATG is the initiation codon; TAG is the stop codon; AATAAA is a polyadenylation signal; AAAAA is the polyA tail; and GGCACAGAGCTGCTCCACAGGCACC is the PCR primer based on Homo sapiens cDNA clone 138761 (Accession No. R63635) similar to the bovine

clone, of Accession No. P08858 neurokinin B precursor used to amplify complete gene.

Figure 4 shows the genomic sequence of neurokinin B, including the 27928  
5 base pair promoter region, the introns, and seven exons (underlined).

Figure 5 shows the results of semi-quantitative PCR for the complete human neurokinin B precursor using mRNA collected at weeks 9, 13 and term. Reverse transcription PCR was performed using mRNA collected at weeks 9,  
10 13 and term (T) to amplify a 733 bp full length neurokinin B precursor cDNA. Primers for  $\beta$ -actin were used as the controls (257 bp). M1 denotes a 1kb DNA ladder; and M2 denotes a 100 bp DNA ladder.

Figure 6 shows HPLC results for oxidised and reduced neurokinin B in  
15 human pregnancy plasma and human term placenta. Placental extracts revealed the peptide to be present in significant amounts ( $21 \text{ pg g}^{-1}$  in early and  $25 \text{ pg g}^{-1}$  in term placenta) and its chromatographic behaviour was identical to synthetic NKB. Partial oxidation of placental NKB during extraction resulted in the production of three oxidised forms in which one or  
20 both of the two-methionine residues were oxidised (**a** in plasma and **b** in placenta). The resulting methionine sulfoxides conferred reduced hydrophobicity, so that they eluted before the reduced form. This elution pattern matched that produced by the partial oxidation of synthetic NKB by hydrogen peroxide. Complete oxidation by hydrogen peroxide resulted in all  
25 the NKB eluting in the position of the first peak. A similar elution pattern was also observed after extraction of NKB from term placenta samples (**b**).

Figure 7 shows the cardiovascular effect of neurokinin B in conscious rats. Changes in blood pressure and heart rate during infusion of saline or  
30 incremental doses of NKB in conscious unrestrained female rats. NKB was infused at doses of  $1.8 \text{ nmol h}^{-1}$  (per kg) from time = 0,  $18 \text{ nmol h}^{-1}$  (per kg) from time = 16 h and  $180 \text{ nmol h}^{-1}$  (per kg) from time = 20 h. Values are mean

± s.e. mean. \* denotes a significant difference from the original baseline and from the values at t = 20 h (Friedman's test).

Figure 8 shows an *in situ* hybridisation of for neurokinin B mRNA in the placenta of humans and rats. **a**, human at term (39 weeks) with human antisense probe **b**, human at term (39 weeks) with human sense probe **c**, rat 18 day placenta with rat antisense probe and **d**, high magnification showing giant cells of the rat placenta expressing neurokinin B. Magnification: **a**, 10x original size, **b** 10x, **c** 16x, **d** 40x.

The present invention is partly based upon the discovery that early and/or excessive release of neurokinin B into the maternal blood stream by the developing placenta can be a cause of pregnancy induced hypertension and pre-eclampsia. In particular, it has been postulated that those likely to suffer from pregnancy induced hypertension or pre-eclampsia have slightly elevated levels of neurokinin B in the maternal blood stream at approximately 10 to 12 weeks into pregnancy. Monitoring of neurokinin B early in pregnancy, for example at 10 to 12 weeks or before, is useful in predicting whether the individual is likely to suffer from pregnancy induced hypertension or pre-eclampsia later in pregnancy, and whether they are likely to suffer from pre-eclampsia related foetal complications such a foetal growth retardation, foetal hypoxia or miscarriage. Measurement of neurokinin B levels after 10 to 12 weeks into pregnancy, for example at 18 weeks may enable the prediction to be confirmed and a diagnosis of pregnancy induced hypertension or pre-eclampsia or related foetal complications to be made. Further, it has been observed that the level of increase in neurokinin B levels after any initial prediction of hypertension or pre-eclampsia correlates with the future severity of the condition. In particular, it has been shown that a relationship exists between the degree of increase in neurokinin B and the future severity of the condition. These observations can be used in the prediction of the future severity of the condition. Also, other post-processing fragments of the human neurokinin B precursor may be involved in the

development of those conditions. In addition, the production of neurokinin B and/or other fragments of human neurokinin B precursor may be associated with the development of hypertension in non-pregnant individuals.

5 In the present invention, foetal complications include any foetal condition which is related to pre-eclampsia. Specifically, foetal complications include foetal growth retardation, foetal hypoxia, pre-term labour, and in severe cases, miscarriage.

10 For the purpose of the present invention, neurokinin B precursor gene products include polynucleotide sequences encoding neurokinin B precursor or neurokinin B, and neurokinin B precursor polypeptides. Polynucleotide sequences include genomic or cDNA sequences, for example those of Figures 3 or 4, and RNA, preferably mRNA. Preferably, the neurokinin B  
15 precursor polypeptides have the sequences shown in Figure 1. Fragments of neurokinin B precursor gene products are fragments which are derived from the precursor gene products and include the polynucleotide or polypeptide sequences encoding neurokinin B, fragments thereof, and other post-processing fragments of the precursor. Preferably the neurokinin B peptide  
20 derived from the precursor has the sequence of Figure 2. Epitopic fragments or variants are those which comprise an amino acid sequence, typically of at least 4 residues, which constitutes a site to which the antibody can bind. A preferred epitopic fragment is the amino acid sequence DMHD of Figure 1.

25 Also included are variants of neurokinin B precursor gene products. Preferably, variants share at least 80%, at least 90%, at least 95%, at least 98% and most preferably at least 99 % sequence identity with the neurokinin B precursor gene products or fragments thereof, and preferably retain the same biological activity as the gene product or fragment.

30

"% identity", as known in the art, is a measure of the relationship between two polypeptide sequences between two polypeptide sequences or two

polynucleotide sequences, as determined by comparing their sequences. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. The alignment of the two sequences is examined and the number of positions giving an exact amino acid or nucleotide correspondence between the two sequences determined, divided by the total length of the alignment and multiplied by 100 to give a % identity figure. This % identity figure may be determined over the whole length of the sequences to be compared, which is particularly suitable for sequences of the same or very similar length and which are highly homologous, or over shorter defined lengths, which is more suitable for sequences of unequal length or which have a lower level of homology.

Methods for comparing the identity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J *et al* , Nucleic Acids Res. 12:387-395, 1984, available from Genetics Computer Group, Madison, Wisconsin, USA), for example the programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (Advances in Applied Mathematics, 2:482-489, 1981) and finds the best single region of similarity between two sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences which are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison, GAP aligns two sequences finding a "maximum similarity" according to the algorithm of Needleman and Wunsch (J. Mol. Biol. 48:443-354, 1970). GAP is more suited to comparing sequences which are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3 for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively. Preferably, % identities and similarities

are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S.F. *et al* , J. Mol. Biol., 215:403-410, 1990, Altschul S.F. *et al* , Nucleic Acids Res., 25:289-3402, 1997, available from the National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)) and FASTA (Pearson W.R. and Lipman D.J., Proc. Nat. Acad. Sci., USA, 85:2444-2448, 1988, available as part of the Wisconsin Sequence Analysis Package). Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S. and Henikoff J.G., Proc. Nat. Acad. Sci., USA, 89:10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide sequences are first translated into amino acid sequences before comparison.

Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a polynucleotide or a polypeptide sequence of the present invention, the query and the reference sequence being optimally aligned and the parameters of the program set at the default value.

The first, second, third and fourth aspects of the invention relate to methods of predicting or diagnosing pregnancy induced hypertension or pre-eclampsia or related foetal complications in a human subject. These methods include, for example, assessing the concentration in a biological sample of neurokinin B precursor gene products, or variants or fragments thereof. These methods preferably comprise comparing the results of an assessment of the concentration of human neurokinin B gene product (e.g. neurokinin B or its precursor) in a sample with expected values or with the values found in the subject at an earlier date.



Preferably these methods are carried out at an early stage of pregnancy, for example at 10-12 weeks for prediction, or 18 weeks for diagnosis.

- 5 These methods may include any means of measuring neurokinin B gene products available to those skilled in the art. Preferably, the methods use the kits of the invention. The methods of the invention comprise at least the step of determining the presence of neurokinin B mRNA, neurokinin B or its precursor, or variants or fragments thereof, in a biological sample; however,  
10 additional steps may also be included. Such additional steps may include one or more of the following: collecting the biological sample; preparing the biological sample; measuring the concentration of target neurokinin B gene products such as polypeptide or polypeptides in the sample; preparing standard curves to predict expected concentrations of the target neurokinin B  
15 gene products in non-pregnant individuals or in pregnant individuals at the same or different stages of pregnancy; comparing the results obtained from a particular biological sample with the appropriate expected values or the appropriate standard curve to determine the severity of the condition; or repeating some or all of the previous steps at a later date to determine if the  
20 severity of the condition has changed.

Suitable methods of detection based on kits will be clear to one skilled in the art and include radioimmunoassay (RIA), enzyme linked immunosorbant assay (ELISA), immunoradiometric assay (IRMA), antisense technology, or  
25 radioreceptor assay (RRA). In the latter, for example the NK<sub>3</sub> receptor or other neurokinin B binding partner may be used in a detection system or biosensor system. Further detection methods may also include as well as radiometric methods, non-radioactive methods such as fluorescence and luminescence.

30

A preferred method is radioimmunoassay, which relies on the interaction of a small amount of radiolabeled peptide, eg neurokinin B, with a limiting amount

of binding partner such as antibody (e.g. specific for NKB). The displacement of radiolabeled peptide by increasing doses of standard peptide is compared to that displaced by unknowns. This is normally monitored by separating binding partner bound label from free label usually by using a precipitation  
5 step which brings down the binding partner followed by centrifugation, although there are adsorbents (e.g. charcoal) which can bind the free labeled fraction and can then be removed by centrifugation. IRMA can be one site or two site and uses an excess of specific binding partner such as antibody which in this case is radiolabeled. In the one site assay, separation is  
10 effected by an excess of peptide linked to a solid phase which removes unreacted binding partner. In the two site method a second specific binding partner (usually linked to a solid phase) is used which is specific to a separate epitope on the peptide. Separation is easily effected by removal of the complex on the solid phase. RRA is similar to RIA in that a limiting  
15 amount of receptor is substituted for the antibody. Often the receptor preparation will be in the form of a membrane preparation so that washing and separation of the bound label can be performed by e.g. centrifugation. The use of enzymes as the signalling moiety in immunometric assays is commonly achieved by cross linking an enzyme to the specific antibody or  
20 the use of e.g. a pig anti mouse antibody cross-linked to an enzyme when a mouse monoclonal antibody is used in the initial reaction.

The above methods may also be used in estimating the likely future degree of pregnancy induced hypertension or pre-eclampsia or related foetal  
25 complications. These methods preferably comprise comparing the results of an assessment of the concentration of human neurokinin B gene product (e.g. neurokinin B or its precursor) in a sample with expected values. It is believed that the tenth week of pregnancy, or later, for example after 18 weeks, may be particularly valuable times at which to assess the presence  
30 (and concentration) of the human neurokinin B gene products.

The methods of the invention are preferably carried out *in vitro*, on a sample removed from the body. Any biological sample may be used in the methods of the invention. Preferred biological samples include blood, saliva or urine.

- 5 The invention also provides a method of preventing or treating pregnancy induced hypertension or pre-eclampsia or related foetal complications in a human subject by the administration of an agent which inhibits the biological effect of neurokinin B. Preferably, such methods are carried out using the kits of the invention. Agents which inhibit the biological effects of neurokinin
- 10 B include any agents that act, for example, by removing the neurokinin B from the plasma; by altering its structure to prevent it binding to receptors; by binding to the receptors directly to block the binding of neurokinin B thereto (but without themselves causing the effects at those receptors normally caused by neurokinin B), by exerting a counter effect to the neurokinin B at
- 15 the same or different receptors or by reducing or preventing gene expression or translation, for example by modulating activity of the neurokinin B gene promoter and/or by using antisense technology. Also included are agents which inhibit the production or processing of the precursor to prevent production of neurokinin B. Within this context, agents inhibiting the
- 20 biological effect of neurokinin B include agents inhibiting the biological effect of any variants or fragments of human neurokinin B or its precursor which are involved in the development of pregnancy induced hypertension or pre-eclampsia or related foetal complications. The principal site of action of human neurokinin B is the NK<sub>3</sub> receptor and therefore preferred agents
- 25 which inhibit the biological effects of neurokinin B for use in the invention include NK<sub>3</sub> receptor antagonists. However, at the high circulatory concentrations found in near term pregnancy, particularly in pregnancy induced hypertensive or pre-eclamptic subjects, neurokinin B may also have significant effects at other receptors (eg the NK<sub>1</sub> or NK<sub>2</sub> receptors) and
- 30 therefore the agents which inhibit the biological effects of neurokinin B for use in the present invention also include agents which prevent neurokinin B's

effects at such other specific receptors, as well as broad spectrum neurokinin antagonists and combinations thereof.

Since 1991, a number of high-affinity nonpeptide antagonists have been reported. Snider R. M., et al., (Science, 251:435 (1991)), and Garret C., et al., (Proc. Natl. Acad. Sci., 88.:10208 (1991)), described CP-96,345 and RP 67580, respectively, as antagonists at the NK<sub>1</sub> receptor, while Advenier C., et al., (Brit. J. Pharmacol., 105:78 (1992)), presented data on SR 48968 showing its high affinity and selectivity for NK<sub>2</sub> receptors. More recently Macleod, et al., (J. Med. Chem., 36:2044 (1993)) have published on a novel series of tryptophan derivatives as NK<sub>1</sub> receptor antagonists. Recently, FK 888, a "dipeptide" with high affinity for the NK<sub>1</sub> receptor was described (Fujii J., et al., Neuropeptide, 22:24 (1992)).

Suitable NK<sub>3</sub> receptor antagonists for use in the present invention include all materials blocking or reducing the effect of neurokinin B at the NK<sub>3</sub> receptor, for example, those materials described in Gao and Peet (Current Medicinal Chemistry, 1999, 6, 375-388), Khavaga and Rogers (Int.J.Biochem Cell Biol. 1996, 28, 7, 721-738), US 5,942,523, US 5,846,973, US 5,491,140, US 5,328,927, US 5,360,820, US 5,344,830, US 5,331,089, US 4,742,156, US 4,665,157, EP 591,040A, WO 94/01402, WO 94/04494, WO 93/011609, Canadian Patent Application 2,154,116, EP 693,489 and Canadian Patent Application 2,151,116. Specific examples of suitable antagonists include the receptor selective ligand, SR 142801 (Edmonds-Alt, et al., Life Sciences, 56:27 (1995)), and the decapeptides of formula: A<sup>1</sup> -D-Pro<sup>2</sup> -His<sup>3</sup> -D<sup>4</sup> -Phe<sup>5</sup> -D-Trp<sup>6</sup> -Val<sup>7</sup> -D-Trp<sup>8</sup> -Leu<sup>9</sup> -Nle<sup>10</sup> -NH<sub>2</sub> wherein A<sup>1</sup> and D<sup>4</sup> are Asp or D-Asp amino acids .

Preferred agents for inhibiting the biological effects of neurokinin B include those which modulate activity of the neurokinin B precursor gene promoter, thus altering the level of transcription of the neurokinin B precursor gene. Examples of such agents include competitive or non-competitive antagonists

of neurokinin precursor B gene promoter transcription factors, agents which inhibit the biological effect of neurokinin B precursor gene promoter transcription factors, agonists of neurokinin B precursor gene promoter inhibitors, and polynucleotide sequences which bind to, and inhibit, neurokinin B precursor gene promoter activity. Preferably, such polynucleotide will be sufficiently complimentary to whole or part of the promoter sequence such that they hybridise thereto and inhibit promoter activity, preferably *in vivo*. Examples of suitable polynucleotide sequences are those which have at least 80%, 85%, 90%, 95%, 97%, 98% and preferably 99% sequence identity with the complement of whole or part of the promoter. Preferably the polynucleotide sequence will be complimentary to a regulatory region of the promoter, for example a transcription factor binding site.

Where the agent is a polynucleotide sequence, it is preferably administered in the form of a vector. The vector may additionally comprise one or more regulatory sequences for activation of expression of the polynucleotide sequence, for example promoters including response elements, consensus sites, methylation sites, locus control regions, post-transcriptional modifications, splice variants, homeoboxes, inducible factors, DNA binding domains, enhancer sequences, initiation codons, and polyA sequences. Such agents may be administered by any suitable gene therapy technique, which will be known to persons skilled in the art.

Administration of pharmaceutical compositions is accomplished by any effective route, e.g. orally or parenterally. Methods of parenteral delivery include topical, intra-arterial, subcutaneous, intramedullary, intravenous, or intranasal administration. Administration can also be effected by amniocentesis related techniques. Oral administration followed by subcutaneous injection would be the preferred routes of uptake; also long acting immobilisations would be used. Also, as the effects of placental NKB will be on peripheral receptors, effectively drugs devoid of side effects to the

central nervous system should be preferably peptide-like in their distribution properties. In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and other compounds that facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of "REMINGTON'S PHARMACEUTICAL SCIENCES" (Maack Publishing Co, Easton PA).

- 10 Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art, in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, etc., suitable for ingestion by the patient.

15 Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable additional compounds, if desired, to obtain tablets or dragee cores.

- 20 Suitable excipients are carbohydrate or protein fillers. These include, but are not limited to sugars, including lactose, sucrose, mannitol, or sorbitol, starch from corn, wheat, rice, potato, or other plants; cellulose such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; and gums including arabic and tragacanth; as well as proteins, such as gelatin and collagen. If desired, disintegrating or solubilising agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

- 30 Dragee cores are provided with suitable coatings such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may

be added to the tablets or dragee coatings for product identification or to characterise the quantity of active compound (i.e. dosage).

Pharmaceutical preparations, which can be used orally, include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with filler or binders such as lactose or starches, lubricants such as talc or magnesium stearate, and, optionally, stabilisers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycol with or without stabilisers.

Pharmaceutical formulations for parenteral administration include aqueous solutions of active compounds. For injection, the pharmaceutical compositions of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilisers or agents, which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner similar to that known in the art (e.g. by means of

conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilising processes). The pharmaceutical compositions may also be modified to provide appropriate release characteristics, e.g. sustained release or targeted release, by  
5 convention means, e.g. coating.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in  
10 aqueous or other protonic solvents that are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilised powder in 1 mM-50 mM histidine, 0.1%-2% sucrose, 2%-7% mannitol at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

15 The agents for use in the invention (eg NK<sub>3</sub> receptor antagonists) can also be modified so that they are only delivered to selected target sites. For example, by adjusting their stability towards proteolytic digestion in the gut or ability not to pass the blood/brain barrier, or by producing composite molecules including a targeting component, e.g. an antibody selective for the  
20 target site.

After pharmaceutical compositions comprising a compound of the invention formulated in an acceptable carrier have been prepared, they can be placed in an appropriate container and labelled for treatment of an indicated  
25 condition. For administration of NK<sub>3</sub> receptor antagonists, such labelling would include amount, frequency and method of administration.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective  
30 amount to achieve the intended purpose. Thus, a therapeutically effective amount is an amount sufficient to ameliorate the symptoms of the disease being treated. The amount actually administered will be dependent upon the



individual to which treatment is to be applied, and will preferably be an optimised amount such that the desired effect is achieved without significant side-effects. The determination of a therapeutically effective dose is well within the capability of those skilled in the art. Of course, the skilled person  
5 will realise that divided and partial doses are also within the scope of the invention.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in any appropriate animal model (eg  
10 primates for pre-eclampsia, rats and guinea pigs for hypertension and other small laboratory animals for use with induced hypertension and induced pre-eclampsia). These assays should take into account receptor activity as well as downstream processing activity. The animal model is also used to achieve a desirable concentration range and route of administration. Such  
15 information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective amount refers to that amount of agent, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity of  
20 such compounds can be determined by standard pharmaceutical procedures, in cell cultures or experimental animals (e.g.  $ED_{50}$ , the dose therapeutically effective in 50% of the population; and  $LD_{50}$ , the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ration  $ED_{50}/LD_{50}$ .  
25 Pharmaceutical compositions, which exhibit large therapeutic indices, are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage varies within this range  
30 depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage is chosen by the individual physician in view of the patient to be treated. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Additional factors, which may be taken into account, include the severity of the disease state. Long acting pharmaceutical compositions might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation. Guidance as to particular dosages and methods of delivery is provided in the literature (see, US Patent No's 4,657,760; 5,206,344 and 5,225,212 herein incorporated by reference).

The agents which inhibit the biological effect of neurokinin B for use in the methods of the invention of preventing or treating pre-eclampsia; or of preparing medicaments for preventing or treating pre-eclampsia; are preferably formulated such that use of the agent is effective in, but not restricted to, the post prandial phase. The agents may for example be selected to be effective over a 24 hour period rather than exclusively in the post-prandial phase. The post-prandial phase is a particularly important time as it is believed that pre-eclampsia is associated with the build-up of toxins in the maternal blood supply due to the failure of the blood to pass through the liver (which normally removes the toxins) because of high pressure in the portal vein. Thus, transient relief of hypertension following meals will allow the blood to pass through the liver at the time when the highest concentration of toxins will be present and will therefore provide a large reduction in the risk of pre-eclampsia whilst producing only a short decrease in the effect caused by the placentally produced neurokinin B. This time limited effect may be achieved by selecting agents with short durations of activity and using appropriate formulations and dosage schedules.

Preferably, methods of prevention or treatment of the conditions addressed herein will begin as soon as possible after the initial prediction or diagnosis is made, for example after 10 weeks into pregnancy. The decision regarding initiation of a course of treatment will of course be the decision of a physician,

and may therefore begin earlier or later. Typically, the course will be given throughout pregnancy or until symptoms subside. This may continue until up to eight weeks after birth. In individuals who have been determined as being at risk of developing foetal conditions such as growth retardation or hypoxia, or pre-eclampsia, (by consideration of other factors such as previous miscarriages or complications in pregnancy) the course may be initiated as soon as pregnancy is confirmed, and may continue until term.

In a further aspect of the invention there is provided the use of a human neurokinin B precursor gene product or a variant or fragment thereof in the manufacture of a diagnostic for use in the prediction or diagnosis of pregnancy included hypertension or pre-eclampsia or related foetal complications. Preferably, the gene product used is neurokinin B, or a variant or fragment thereof, for example in the production of a diagnostic comprising a binding partner specific for neurokinin B. Preferably, the variants or fragments are epitopic. It is envisaged that other gene products could also be used, for example regulatory sequences of the neurokinin B precursor genomic sequence, or neurokinin B precursor mRNA in the production of antisense sequences.

The polypeptides used include human neurokinin B or its precursor, or variants or fragments thereof. Preferably, the polypeptides comprise the sequence of Figure 1 or Figure 2 respectively. Preferably, the fragments or variants are epitopic, as defined above.

These polypeptides may be produced in isolated, substantially pure form or as recombinant polypeptides. Method for doing so will be clear to one skilled in the art. These will include, for example, recombinant techniques or extraction, gel separation or more commonly, for peptides the size of neurokinin B, chemical synthesis, eg liquid and solid phase peptide.

In a further aspect of the invention there is provided the use of an agent which inhibits the biological effect of neurokinin B in the manufacture of a medicament for the prevention or treatment of pregnancy induced hypertension or pre-eclampsia or related foetal complications. Preferably,  
5 the agents are those defined above.

In a further aspect of the present invention there are provided kits for the predicting the onset of, diagnosing, or estimating the future severity of pregnancy induced hypertension or pre-eclampsia or related foetal  
10 complications. The kits of the invention comprise a means for detecting the production of human neurokinin B gene products such as polynucleotides or polypeptides encoding neurokinin B or its precursor, or fragments or variants thereof, by the subject. Thus the kits will commonly comprise one or more of:  
15 a binding partner to neurokinin B or its precursor; neurokinin B polypeptide or variants or fragments thereof; and/or polynucleotide sequences which hybridise to a sequence encoding neurokinin B or a variant or fragment thereof.

By binding partner is meant any substance capable of detecting (and binding  
20 to) the target, eg an antibody. Preferred binding partners for use in the kits of the invention are antibodies which are specific for neurokinin B precursor, or epitopic fragments or epitopic variants thereof. Preferred are antibodies to neurokinin B and antibodies to the human neurokinin B precursor. Most preferred are antibodies which are specific for neurokinin B, but antibodies  
25 specific to any other breakdown products of the neurokinin B precursor which remain in the body for a measurable time may also be used. These antibodies are capable of binding fragments of the human neurokinin B precursor to identify the production of the precursor by the human body. The antibodies of the invention may be, for example, polyclonal, monoclonal,  
30 chimeric or humanised antibodies or fragments thereof. Binding partners which cross react with related peptides such as Substance P or NKA, for

example, may be useful as a medicament or in diagnosis, as they share a common sequence (FVGLM-NH<sub>2</sub>) with neurokinin B.

5 Methods of producing such antibodies will be apparent to one skilled in the art. For example, in the case of polyclonal antibodies, by standard methods of animal immunisation or, for monoclonal antibodies, by the well-known methods of Köhler and Milstein, or by use of the methods discussed in US 5,844,080. Chimeric antibodies can be made by genetic engineering techniques, and are antibodies in which the constant region is human in  
10 origin, but the variable regions are derived from, for example, a mouse antibody. The advantage of chimeric antibodies is to reduce immunogenicity. Humanised antibodies take this principle even further, in that only the complementarity determining regions and a minimum number of further amino acids in the variable regions are derived from an animal such as a  
15 mouse. The rest of the antibody structure is human in sequence, and is recognised by the human immune system as human (see, for example, Queen et al, PNAS, USA 86 (December 1989), 10029-10033).

Polynucleotides of the kits of the invention are preferably those which  
20 hybridise to a sequence encoding neurokinin B or its precursor, or a variant or fragment thereof, or complements thereof, under stringent conditions. Preferred are polynucleotide sequences which hybridise to the nucleotide sequence of Figure 3 or Figure 4, or their complements, under stringent hybridisation conditions. Stringent conditions are, for example, 6x SSC at  
25 65°C. Preferably, such polynucleotide sequences have at least 85%, and least 90%, at least 95%, preferably at least 98% and most preferably at least 99% sequence identity with the complement of the reference sequence. Such polynucleotide sequences are preferably at least 10 nucleotides in length, and will be useful in detecting expression of neurokinin B or its precursor.  
30 Such polynucleotides are useful in antisense technology or diagnostic PCR.

Means of producing the polynucleotides of the invention will be clear to those skilled in the art, for example, they may be produced synthetically or by probing an appropriate cDNA or genomic library (particularly a placental cDNA library).

5

The kits of the invention may also comprise instructions for the performance of an assay for predicting or diagnosing the levels of neurokinin B in a biological sample (this may either be by direct measurement of neurokinin B or by measuring the concentration of human neurokinin B precursor, or a  
10 fragment thereof, and using this value to predict the amount of neurokinin B present). The components of the commercial neurokinin B radioimmunoassay kit RIK 7357 by Peninsula Laboratories, Belmont, CA, USA can be used in the present invention. The kits of the invention preferably also comprise a key, showing the correlation between the levels of  
15 neurokinin B gene product in the biological sample and diagnosis of pregnancy induced hypertension or pre-eclampsia or related foetal complications, and/or the likely future onset and/or severity of these conditions.

20 Also provided are kits for the prevention or treatment of pregnancy induced hypertension or pre-eclampsia or related foetal complications, comprising means for inhibiting the biological effect of neurokinin B or its precursor in a subject. Preferably, such means include those agents defined above. In particular, the antibodies or polynucleotide sequences as described above  
25 may also be useful in these kits for inhibiting the biological effect of neurokinin B or its precursor. The kits preferably also contain instructions for use of the kit to prevent or treat pregnancy induced hypertension or pre-eclampsia or related foetal complications and/or a key showing the correlation between the amount of agent used and the likely effect on the  
30 condition.

Pre-eclampsia may also be alleviated by modifying the diet of a human subject to reduce the content of toxins (e.g. alkaloids) and toxin generating substances therein. Toxin generating substances include proteins which are  
5 digested in, and absorbed from, the gut as amino acids most of which are toxic if they circulate in blood in too high concentrations. Normally any amino acids in excess of daily requirement are immediately deaminated by the liver and metabolised. Increasing the proportion of carbohydrates in the diet may also be of particular benefit. The dietary pattern of the subject may also be  
10 modified to prevent peak concentrations of potential toxins appearing in the portal vein, for example by substantially reducing the size of individual meals (and increasing the frequency of small meals).

Agonists of neurokinin B may also be used as pharmaceutical agents where  
15 an increase in blood pressure or decrease in blood volume is considered to be beneficial. Suitable agonists include any acting to supplement or mimic the effect of neurokinin B at the NK<sub>3</sub> receptor (or at any other receptor), for example senktide or [MePhe<sup>7</sup>] NKB.

20 The present invention also provides means of screening potential effective agents (eg NK<sub>3</sub> receptor antagonists and agonists) by testing their ability to block (or enhance) the hypertensive effect of neurokinin B in an appropriate model. Once suitable agents have been identified, they may then further be tested to determine their potential in preventing or treating hypertension;  
25 pregnancy induced hypertension or pre-eclampsia, and used accordingly. All agents identified by such a process (other than presently known materials) are included in the present invention. Screening methods include large array techniques such as the Vilsips™ technology of Affymetrix Inc; see, eg, EPB No. 0476014.

30

Transfected cells lines containing the cloned NK<sub>3</sub> (or NK<sub>1</sub> or NK<sub>2</sub>) receptor could be used in receptor binding and cell signalling pathway studies in a

way clear to one skilled in the art. Essentially, either cells lines expressing endogenously high levels of neurokinin receptors or cell lines transfected with cloned cDNA constructs of the neurokinin receptor may be used to produce membrane preparations. Membrane preparations, of purified receptors in solution or after reconstitution into phospholipid membranes, may then be used to assess receptor binding with labelled agonists and/or antagonists of neurokinin B. The effects of the action of the agonists and antagonists can be assessed using standard cell signalling assays. These will be typical of those routinely performed when using G-protein coupled receptors systems in a way clear to one skilled in the art (including such assays as receptor binding, cyclic AMP determination, protein kinase C, inositol triphosphate concentrations etc.). These studies could also be performed in animal models including the guinea pig and rat chronically infused with agonist to determine the long and short-term effects of neurokinin B, neurokinin B agonists and neurokinin B antagonists. Effects such as changes in heart rate, blood pressure, blood volume and weight of internal organs (e.g. uterus, placenta) may be measured.

## **EXAMPLES**

### **Example 1**

#### **Production of human neurokinin B precursor cDNA**

The cloning of placental cDNA, using the following methods, was used to identify the human neurokinin B precursor having the polypeptide sequence shown in Figure 1. The peptide sequence of neurokinin B in the precursor is underlined (the C-terminal G residue ends up as the amide on the C-terminal M in the final processed peptide of Figure 2). The cloned placental cDNA of the human neurokinin B precursor is shown in Figure 3 and has (underlined) the ATG initiation codon at 26-28, the TAG stop codon at 389-391, the AATAAA polyadenylation signal at 659-663 and the polyA tail starting at 680.



Human placental tissue was obtained from pregnancy terminations at weeks 9 and 13 of gestation and term. Samples were collected in compliance with and approval from the Local Research Ethics Committee. RNA was extracted essentially as described by Chomczynski, P. and Sacchi, N. (1987) Analytical Biochemistry, 162, 156-159.

The full-length preproneurokinin B precursor was amplified using RT-PCR from total human term placental RNA. This was done using the SMART RACE cDNA amplification method (Chenchik, A. *et al* (1998)). In RT-PCR Methods for Gene Cloning and Analysis. Eds. Siebert, P. and Larrick, J. (BioTechniques Books, MA), 305-319). Essentially, after total RNA extraction, reverse transcription was performed using a cDNA synthesis primer (5'AAGCAGTGGTAACAACGCAGAGTAC(T)<sub>30</sub>N<sub>1</sub>N3') which contained a 3' anchor sequence. 3' race was performed using a 5' gene specific primer (5'GGCACAGAGCTGCTCCACAGGCACCAT 3') derived from the Homo sapiens cDNA clone 138761 similar to bovine P08858 neurokinin B precursor. The resulting PCR fragment was gel purified following gel electrophoresis and cloned into the expression vector pGEM-T Easy. The resulting clones were sequenced and compared to submitted sequences in the GenBank database using the BLAST program (Altschul, S.F., *et al* (1990) J.Mol.Biol. 215:403-410).

## **Example 2**

### **Semi-Quantitative PCR to measure NKB in placenta**

Semi-quantitative PCR as described below was used to measure the mRNA expression of neurokinin B in placenta collected at 9 weeks, 13 weeks and at term. This showed differences in a degree of expression between the first trimester and term placenta. Expression levels were up by five times at term, as shown in Figure 5.

30

SMART RACE placental cDNA was amplified using a 5' gene specific primer (5'GGCACAGAGCTGCTCCACAGGCACCAT 3') derived from the Homo

sapiens cDNA clone 138761 similar to bovine P08858 neurokinin B precursor and a 3' SMART anchor sequence primer. A specified primer pair for  $\beta$ -actin was used for normalisation. PCRs were performed using twenty-one cycles of 95°C for 30 sec and 68°C for 2 min. The primers were chosen deliberately to have high annealing temperatures so that the PCR reactions could be performed two step to reduce the possibility of non-specific products being formed. The number of cycles required to obtain a reproducible exponential amplification of the  $\beta$ -actin RT-PCR product was determined by terminating control reactions at 15, 18, 21, 24 and 30 cycles respectively. These experiments were used to check the accuracy, efficiency and amount of total RNA needed to obtain a semi-quantitative amplification in order to optimise the levels of  $\beta$ -actin PCR product produced. The PCR products were visualised by UV illumination following electrophoresis (A 1kb DNA ladder (M1) and 100bp DNA ladder (M2) are shown in Figure 5 also).

### **Example 3**

#### **Neurokinin B extraction from placental tissue and plasma**

Testing of placental extracts using the techniques set out below revealed neurokinin B to be present in significant amounts and its chromatographic properties in HPLC were identical to synthetic neurokinin B. It also displayed the same degree of loss of hydrophobicity (on HPLC) after oxidising its methionine residues. Oxidisation was found to give three peaks of double oxidised (1), single oxidised (2) and non-oxidised forms (3), see Figure 6. Figure 6(a) shows oxidised and reduced neurokinin B separated by RPHPLC from human pregnancy plasma and Figure 6(b) shows separation of condensed and reduced neurokinin by RPHPLC extracted from human term placenta.

#### **Extraction of neurokinin B from placenta**

Whole placentae were weighed and washed immediately after delivery with 150 mM sodium chloride solution containing 10 mM EDTA at pH 7.5. A

tissue sample not exceeding 100g was excised and homogenised in 100 ml saline/EDTA solution using a blender with a glass vessel. Protease inhibitors, phenylmethanesulphonylfluoride, N-ethylmaleimide, and pepstatin were added from a stock solution in methanol. After 20 seconds 800 ml of methanol were added and blending was continued for a further minute. The mixture was decanted into 200 ml polypropylene centrifuge tubes and subjected to centrifugation at 4°C and 3000 X g for 30 minutes. The supernatant was separated and stored overnight at 4°C resulting in further precipitation that was removed by centrifugation. The volume of each extract was reduced to less than one eighth of the initial volume and then diluted by addition of three volumes of water containing 0.1% trifluoroacetic acid (TFA). Any trace of suspended matter was removed by a final centrifugation step. The volume of extract was recorded and an amount corresponding to 20g of placenta reserved for solid phase extraction using Sep-Pak C18 3CC cartridges (Waters Chromatography Division, Millipore Corporation, Milford, MA, U.S.A.). Cartridges were primed prior to use by perfusion with 2 ml of the following solutions; 1) water containing 0.1% TFA and 0.1% Polypep gelatine hydrolysate (Sigma-Aldrich, Poole, UK), 2) water containing 0.1% TFA, 3) water containing 80% v/v acetonitrile and 4) water containing 0.1% TFA. Each extract was passed through a prepared cartridge, which was then washed with 2 ml 0.1% TFA in water, 2 ml 0.1% TFA in water containing acetonitrile 10% and 20% TFA. The column was eluted with 2 ml of 30%, 40% and 50% acetonitrile in water containing 0.1% TFA. Eluted fractions were reduced to dryness under vacuum after adding 1 mg of mannitol and 100 µg Polypep. Smaller placentae obtained from abortions were treated as above but dissociated in a glass homogeniser retaining the same proportions of buffer and methanol to placental weight.

#### **Extraction of neurokinin B from plasma**

Neurokinin B standards were prepared in pooled plasma from the blood of five young males taken into EDTA. The standards contained 1280, 640, 320,

160 and 80 pg/ml neurokinin B. Each 2ml of sample of plasma standard was acidified by addition of 220  $\mu$ l 1M HCl containing 0.21M glycine. They were then diluted to 10 ml with 0.9% saline and subjected to centrifugation at 3000 X g for 20 minutes to ensure complete clarity. Sep-Pak C18 1CC cartridges were primed as described above for Sep-Pak C18 3CC cartridges. After loading, cartridges were washed with 1 ml 0.1 M HCl containing 0.02M glycine followed by 1 ml 0.1% TFA in water. Further washes with 1ml 0.1% TFA in water containing 10 and 20% acetonitrile were followed by elution with 1 ml 0.1% TFA in a mixture of 50% water and acetonitrile. Eluted fractions were reduced to dryness under vacuum after adding 1 mg of mannitol and 100  $\mu$ g Polypep. The acidification step ensured that we were extracting already processed mature peptide as it is possible that inactive circulating precursor could be cleaved by endogenous plasma proteases to produce immunoreactive peptides unless precautions are taken.

#### **Example 4**

##### **Measurement of NKB in placental tissues and plasma**

Placental and plasma extracts were reconstituted in 500  $\mu$ l of buffer supplied as part of a commercial neurokinin B radioimmunoassay kit RIK 7357 by Peninsula Laboratories, Belmont, CA, USA to which had added 0.2% Igepal CA-630 non-ionic detergent (Sigma). Sub-samples of 25  $\mu$ l were taken from extracted and non-extracted standards and mixed with 75  $\mu$ l of the above buffer. Standards were prepared in buffer containing Igepal, but to which had been added 200  $\mu$ g/ml Polypep. Anti-neurokinin antibody solution (100  $\mu$ l) was added to all assay tubes except blanks and the assay was conducted as described in the "General Protocol for Radioimmunoassay Kit" instructions. Assays were performed in duplicate and results were corrected with reference to extracted standards.

The plasma and placental levels of neurokinin B in various human volunteers and rats were measured by the above methods. The results of the plasma samples are summarised in Table 1. Placental samples were collected from weeks 7 to 15 of pregnancy, and all seven were shown to contain equivalent  
5 significant amounts of neurokinin B; however concentrations of plasma NKB detected at term were in the 100 picomolar range that would be expected to have effects on the maternal cardiovascular system. Plasma samples taken from non-pregnant volunteers all had low levels of the peptide, as did the majority of plasma samples taken from individuals who had been admitted for elective  
10 abortions at weeks 7 to 15. Four samples from this latter group had concentrations equivalent to those found at term. This suggests that the placenta from this individual may have started to secrete supra-physiological concentrations of neurokinin B early in pregnancy. Samples of patients in late pregnancy suffering from hypertension and pre-eclampsia all had  
15 concentrations in the nanomolar range suggesting that raised neurokinin B may be responsible for their symptoms.

Table 1

Week of Pregnancy	Nmol/l NKB in normotensive pregnancies
6	0
9	0
9	0.97
10	0.535
13	0
13	0
13	0.083
13	0.511
14	0
14	0
14	0.511
17	0.182
17	0.182
18	0
23	0.12
24	0
25	0.17
27	0
28	0
28	0.033
31	0
31	0.031
32	0
33	0
37	0
38	0.07
39	0.138
40	0.05
40	0.2
41	0.118

Table 2

Week of pregnancy	Nmol/l NKB in pre-eclamptic pregnancies
30	3.964
34	6.156
36	3.796
37	2.141
38	2.752
39	2.004
39	6.288
39	0.98

5 Table 3

Patient number	Nmol/l NKB in normotensive pregnancies at term
1	0
2	0
3	0
4	0
5	0
6	0
7	0
8	0.084
9	0.118
10	0.143
11	0.22
12	0.226
13	0.228
14	0.398
15	0.521
16	1.317

## CLAIMS:

1. A kit for the prediction or diagnosis of pregnancy induced  
5 hypertension, pre-eclampsia or related foetal complications comprising a binding partner, eg an antibody, to neurokinin B precursor gene product or variant or fragment thereof.
2. A kit according to claim 1 further comprising instructions for the  
10 performance of an assay for predicting the levels of neurokinin B in a biological sample and correlating the assay results with the likely future development of pregnancy induced hypertension or pre-eclampsia or related foetal complications respectively.
- 15 3. A kit for use in estimating the likely future degree of pregnancy induced hypertension or pre-eclampsia or related foetal complications, comprising a binding partner, eg an antibody, to neurokinin B precursor gene product or variant or fragment thereof, together with instructions for the  
20 performance of an assay for predicting the levels of neurokinin B in a biological sample and correlating the assay results with the predicted future severity of pregnancy induced hypertension or pre-eclampsia or related foetal complications, respectively.
4. A kit as claimed in any one of claims 1 to 3 wherein the binding partner  
25 is an antibody specific for neurokinin B precursor, or neurokinin B or an epitopic fragment or epitopic variant thereof.
5. A kit according to any one of claims 1 to 4 wherein the binding partner  
30 is an antibody specific for the human neurokinin B precursor having the sequence of figure 1 or an epitopic variant or epitopic fragment thereof.



6. A kit as claimed in any one of claims 1 to 5 which is a radioimmunoassay kit, an enzyme linked immunosorbant assay kit, an immunoradiometric assay kit or a radioreceptor assay kit.

5 7. A method of preventing or treating pregnancy induced hypertension or pre-eclampsia or related foetal complications in a human subject by the administration of an agent which inhibits the biological effect of neurokinin B.

8. The method as claimed in claim 7 wherein the agent which inhibits the  
10 biological effect of neurokinin B is an NK<sub>1</sub>, NK<sub>2</sub> or NK<sub>3</sub> antagonist.

9. The method as claimed in claim 8 wherein the NK<sub>3</sub> antagonist is a decapeptide with the following formula : A<sup>1</sup> -D-Pro<sup>2</sup> -His<sup>3</sup> -D<sup>4</sup> -Phe<sup>5</sup> -D-Trp<sup>6</sup> -  
Val<sup>7</sup> -D-Trp<sup>8</sup> -Leu<sup>9</sup> -Nle<sup>10</sup> -NH<sub>2</sub> wherein A<sup>1</sup> and D<sup>4</sup> are Asp or D-Asp amino  
15 acids or SR 142801.

10. The method as claimed in claim 7 wherein the agent which inhibits the biological effect of neurokinin B is one which modulates the activity of the neurokinin B precursor gene promoter.  
20

11. The method as claimed in any one of claims 7 to 10 wherein the agent is selected and administered such that it effective over a 24 hour period.

12. Use of human neurokinin B precursor gene product or variant or a  
25 fragment thereof in the manufacture of a diagnostic for use in the prediction or diagnosis of pregnancy-induced hypertension or pre-eclampsia or related foetal complications.

13. Use of a human neurokinin B precursor gene product or variant or  
30 fragment thereof according to claim 12, wherein the gene product is human neurokinin B precursor or human neurokinin B, or an epitopic variant or epitopic fragment thereof.

14. Use of neurokinin B in the manufacture of a diagnostic for use in the prediction or diagnosis of pregnancy induced hypertension or the diagnosis of pre-eclampsia or related foetal complications, according to claims 12 or 13.

15. Use of an agent which inhibits the biological effect of neurokinin B in the manufacture of a medicament for the prevention or treatment of pregnancy induced hypertension or pre-eclampsia or related foetal complications.

16. The use as claimed in claim 15 wherein the agent which inhibits the biological effect of neurokinin B is an NK<sub>1</sub>, NK<sub>2</sub> or NK<sub>3</sub> antagonist.

17. The use as claimed in claim 15 wherein the NK<sub>3</sub> antagonist is SR 142801, or the decapeptides with the following formula : A<sup>1</sup> -D-Pro<sup>2</sup> -His<sup>3</sup> -D<sup>4</sup> -Phe<sup>5</sup> -D-Trp<sup>6</sup> -Val<sup>7</sup> -D-Trp<sup>8</sup> -Leu<sup>9</sup> -Nle<sup>10</sup> -NH<sub>2</sub> wherein A<sup>1</sup> and D<sup>4</sup> are Asp or D-Asp amino acids.

18. The use as claimed in claim 15 wherein the agent which inhibits the biological effect of neurokinin B is one which modulates activity of the neurokinin B gene promoter.

19. The use as claimed in any one of claims 15 to 18 wherein the medicament is formulated such that the agent is effective over a 24 hour period.

20. A method of predicting or diagnosing pregnancy induced hypertension or pre-eclampsia or related foetal complications at an early stage in a human subject by assessing the concentration in a biological sample, e.g. blood, of human neurokinin B precursor gene product or a variant or a fragment thereof.

21. A method of predicting or diagnosing pregnancy induced hypertension at an early stage in a human subject or of predicting pre-eclampsia or related foetal complications at an early stage in a human  
5 subject by assessing the concentration in a biological sample, e.g. blood, of neurokinin B or its precursor.

22. A method according to claim 21 wherein neurokinin B and its precursor have the sequences of figures 1 and 2 respectively.

10

23. The method as claimed in claims 20 to 22 comprising the use of a kit as defined in any one of claims 1 or 2.

24. A method of estimating the likely future degree of pregnancy induced hypertension or pre-eclampsia or related foetal complications in a human  
15 subject by assessing the concentration in a biological sample, eg blood, of human neurokinin B precursor gene product or a variant or a fragment thereof, and correlating the result with the predicted future severity of pregnancy induced hypertension ore pre-eclampsia or related foetal  
20 complications.

25. A method according to claim 24 comprising assessing the concentration in a biological sample, e.g. blood, of nuerokinin B.

25 26. The method as claimed in any one of claims 24 or 25 comprising the use of a kit as defined in any one of claims 3 to 5.

27. The method as claimed in claim 26 wherein the kit comprises an antibody specific for neurokinin B.

30

28. The method as claimed in claim 26 or claim 27 wherein the kit is a radioimmunoassay kit, an enzyme linked immunosorbant assay kit, an immunoradiometric assay kit or a radioreceptor assay kit.

5 29. The use of neurokinin B or an agonist thereof in the reduction of blood volume in cases of hypotension.

30. The use of neurokinin B or an agonist thereof in the preparation of a medicament for the reduction of blood volume in cases of hypotension.

10

31. A method of alleviating pre-eclampsia in a human subject by modifying the diet of the human subject to reduce the content of toxin generating substances therein.

15 32. A method of alleviating pre-eclampsia in a human subject including modifying the dietary pattern of the subject to reduce concentrations of potential toxins in the portal vein.

20 33. A dietary methodology for the alleviation of pre-eclampsia in a human subject in which the amount of toxin generating substances is reduced.

1 / 20

## FIG. 1

THE AMINO ACID RESIDUE SEQUENCE OF THE HUMAN NEUROKININ B PRECURSOR

MRIMLLFTAILAFSLAQSF~~GAVCKEPQEEVVP~~GGGRSKRDPDLYQLLQRLFKSHSSLEGLLKALSQASTDPK  
 ESTSPEKRDDMHDFVGLMGKRSVQPDSPD~~VNQENV~~PSFGILKYPPRAE

## FIG. 2

THE AMINO ACID SEQUENCE OF NEUROKININ PEPTIDE

DMHDFVGLM-NH<sub>2</sub>

## FIG. 3

THE CLONED FULL-LENGTH PLACENTAL cDNA OF THE HUMAN NEUROKININ B PRECURSOR

GGCACAGAGC	TGCTCCACAG	GCACCATGAG	GATCATGCTG	CTATTCACAG	50
CCATCCTGGC	CTTCAGCCTA	GCTCAGAGCT	TTGGGGCTGT	CTGTAAGGAG	100
CCACAGGAGG	AGGTGGTTCC	TGGCGGGGGC	CGCAGCAAGA	GGGATCCAGA	150
TCTCTACCAG	CTGCTCCAGA	GACTCTTCAA	AAGCCACTCA	TCTCTGGAGG	200
GATTGCTCAA	AGCCCTGAGC	CAGGCTAGCA	CAGATCCTAA	GGAATCAACA	250
TCTCCCGAGA	AACGTCACAT	GCATGACTTC	TTTGTGGGAC	TTATGGGCAA	300
GAGGAGCGTC	CAGCCAGACT	CTCCTACGGA	TGTGAATCAA	GAGAACGTCC	350
CCAGCTTTGG	CATCCTCAAG	TATCCCCCGA	GAGCAGAATA	GGTACTCCAC	400
TTCCGGACTC	CTGGACTGCA	TTAGGAAGAC	CTCTTTCCCT	GTCCCAATCC	450
CCAGGTGCGC	ACGCTCCTGT	TACCCTTTCT	CTTCCCTGTT	CTTGTAACAT	500
TCTTGCTGCTT	TGACTCCTTC	TCCATCTTTT	CTACCTGACC	CTGGTGTGGA	550
AACTGCATAG	TGAATATCCC	CAACCCCAAT	GGGCATTGAC	TGTAGAATAC	600
CCTAGAGTTC	CTGTAGTGTC	CTACATTAAA	AATATAATGT	CTCTCTCTAT	650
TCCTCAACAA	<u>TAAAGGATTT</u>	TTGCATACGA	<u>AAAAAAAAAA</u>	<u>AAAAAAAAAA</u>	700
<u>AAAAAA</u>					706

2 / 20

## FIG. 4

```
1 AGGCTACTGT AGGTAACCAC CCAGCTTGGT TCTTCAGCTC CACATGGTGG GGTTAGGAGA
61 GGAGGAGGAG GGAGATGGAT GGAACCAATT AGGAACAGCA CCTGGGCTCC TCACAGGAAT
121 GAACCAAGTCA TGCCATTTGC ATGTAAACAG CTTCCCACTT CTCTCCTCAT CCTACCAAAT
181 GCTCCCAACC CTGGGTCTCTG GCCCATGTTC TTTGCCCACA CAGCCCTGTA ATTAGCTGGG
241 TAATGAGAAG CTTTAAATGA GTCCCATTAG CATCTCGTGT AATAAAGAGG CCTTGAGACC
301 CAGCTGCTGT CCTCACTTTG GGATGAACAC GGGTCCCTGT GTAGCCAGTG ACTTCTGTCA
361 GTACAGTCTA AGTTCTCGGA TGGGGTGGGA GACAAACATT TCAGGACCCC AGCAGCACTT
421 GAGAGGTTCC ATGGTGGATC CATGTTTTTG ACTGTGATAC AAGAACTTG GCTCTGGCTT
481 CCTTGTTTCAT TTTGTAAATA ACATTTTTTC TTCTTTAAG AGACAGAGTC TTACTTTGTT
541 GCCCAGGCTG GAGTGTAGCA ATGCAATTAT AGCTCACTGC AGCCTCAACC TCCTGGGCTC
601 AAGTGATCCT CCTGCCTCAG CCTCTGGGAT AGCTGGGGCC ACAGGCATGC ACCACCATGC
661 CTGGCTAATT TTTAAAAATG TTTTGTAGA GATGGGGTCT TACTTGCTAT GTTGCTCAGA
721 CTGGTCTCGA ACTTCTGGCT TCAAGCAATT CTCCCACCTC GCCCTCCTAA AGTGCTGGGA
781 GTATGGGCAT GAGCCACCAT GTCCAGCCTT GTAAATACAT TTTTATTGAG CACCTATTAT
841 ATGTCAAACA TTATAAAGTG AGGGATACAG TAGCAAACAA AACAGACAAA AATTTTTGCC
901 ATCATGACAC TTATATTCCT GGGTGGGAGT GGTGATAGAA AGACAATAAG TAAAATACTT
961 AGCATAGTGG ATGTAATAAG TTCATGAAGG GAAAAATGGG AGTGAGGTAT ATGGAATTTT
1021 GGGGTGGTGA TAATTTTAAA TAGGGTGATT GGGGAATGCT TTGTTGCACA GATTGTTTTT
1081 GTAGTAAATA TGAGATAAAG ATACGGTTCT CTCCCAAAC CAAAATGTAG AAGAGTAGAA
1141 GGTCCTCAAT CTTCAAGTCT CTTGGAGAGG GGGGCCACCC ATTCCGTCTG GGACAGTTAA
1201 CTGTTCCCTC ACAGGTCAAA GTTTATGCCA GTGCAGTAAA AAGAGTGGGA GACCTGGGGT
1261 GAGACAAAACC TGGATTTGAG GCTGTTCTTC ACTGATTAGT AGCCATATGT ACTGGAGCAA
1321 GTGACTGAAC CTTCTGAGCC TGTTTTCTCA TCTGGAAAAT CAGAATATTT CCTACTTACA
1381 TGGTCATGGT GATGAAAACC AGATGGACTG CTCCATGCCA AAGCACCTG CAAACATTCA
1441 AACCTGCAC CCATTACAAA TACTGGGCTG ACGGATGGCT CTGGCTTTGC TTTTGCATCT
1501 CCGCTGTCTC ATTCAGCAGC AGCATCTGGC TCTGGCTCTC GGCTCTGATC CTGGTTCTGA
1561 CTCTCCCCTG GAGCTCTCTC CCTTGGGTGA GAAATAAGCA GATAATCTCC CTCATCTGTG
1621 TGTGGTGTGA ACAAGAGGCT TGAAAGGTCA GAGAAGAAGA TGCCTGAACT GCAGGGAGAC
1681 AGATTAGAGT GGGGAAAATG TAACTCTGAG GAAAAAGGGA AGCAATTAAG AGATCAAGGC
1741 CAGGGGCAGT GGCTCATGCC TGTAATCCCA ACACCTTGGG AGGCTGAGGC GGGCAGACCA
1801 TGAGGTCAGG AGTTCGAGAC CAGTCTGGCC AACATAGTGA AACCCCGTCT CTACTAAAAA
1861 TACAAAAAAA TTAGCCAGGT ATGGTGGTGT GCACCTGTAA TCCCAGCTAC TTGGGAGGCT
```

3 / 20

```
1921 GAGGCAGAAG AATTGCATGA ACCCGGGAGG CAGAGGTTGC GGTGAGCCGA GATTGAACCA
1981 TTGCACTCCA ACCTGGGCAA CAGTGTGAGA CTCTGTCTCC AAAAAAAAAA AAAAAAAAAA
2041 AAATCAAGGC CGGGGAGGGG GCAGGGGTGG CACAGCTATC GAGTTCTGTT CATCCTCTGT
2101 GAGATTACAT CAGGAGGTGT AAAAGAACTC TAGAAGAATG AAGCTAAGTC CAGCTGATTC
2161 AGGGTTCAAG AAGGATTGAG GTGGGAGAGG CATCATGACC ACTGGTGAGG AGTGGAGGAA
2221 GGCCGACACT GGAGCTTTCT TTGCCCAAGC AGAGGAGGGG TGTGACACTC TTGAGGACCA
2281 ATGTAATGGC GCAGCTCCCT CTGGGAGGGG GAAAGGAGAG GACTGGAGGG GATGCTAAAC
2341 TGACCTTCTA ACCTTCAGGG GCCTGAGTCT GGTGTCTCTG GGTGGGGAGG GGCCTGCTCC
2401 TGAAACTGTT TTAGCCCAGA AGTCAGGCCT GAAGGTTAAA GGGCAAGGAG CTGGTGGATG
2461 AACAAGGTGG GGAAAGAGGC CCAGGGTCCA CATCTACTGA GCTGGACTCA GGCATGGGAA
2521 TTGGTGTGTT GAGGGCCAAG ACACTTGGCC TCCTAAAAGT TTGCTGAAAA TCACTGACAT
2581 GAGAGTAATT GATTTATAGG AGAAAAGGTA GATAAATTTA TTTAATATGT ATATATGAGC
2641 ACCTTTAGAA TGAAGACCCA AAGATATAGG GGAAATTGCC AGTTATTTAT TTATTTTTTT
2701 TGGAGATGGA GTCTCACTGT GTCTGCCAGG CTAGAGTGCA GTGGCATGAT CTCGGCTCAC
2761 TGCAACCTCC GCCTGCTGGG TTCAAGCAAT TCTCCTGCCT CATCCTCCTG AGCAGCTGTG
2821 ACTACAGGCA CGCACCACCA TGCCCGGCTA ATTTTTTGTA TTTTTTAGTA GAGACAGGGT
2881 TTCACCATGC TGGCCAGGCT GGTCTGGAAC TCCTGACCTT GTGATCCGCC CGCCTTGGCC
2941 TCCCAGAGTG CTGGGATTAT AGGCATGAGC CACCGCCCCC AGCCTGAAAT CGCCAATTTT
3001 ATGTTTATGT TTTACAAAGT ATGGACAGCT GTGTAGAAAT ATGACTGGAC AGAAGGGCAT
3061 GCTCTAATGT TAACAGACTG AGTGGGGAAA CCCAGGAAGG CCTGTTGAGA TTCCTCCTGG
3121 CCTCTCTCAT TCCTTCCTTC TGGGTATGGG GCAGGACCCT CTCTGGAATG GGGAGATCTT
3181 AGGACCTAAG TTAAATAAGG TAGGTCAGAT AATTTTTTAT GGCCAGTTTT TACATACAGT
3241 AATTTTAGGT TTTATGGCTG GCTTTGGGGA AAAGAGGTCC TGGTTTTTAT AGCTGGCCTT
3301 GGGGGAGAAT GGGACCCAGC AACAGGAGGA CAGGAGAGGG TCAGAGAAAA ACTTCTGCTT
3361 CTGAGGCTGC TACTGAGGCC TTCATTTTAG GGTATTGTCT TCTGAGCCCC AGCATTCCCTC
3421 GGTGTGAAAA ATTTTAAAGA AATTTTATAG TCCAGAAATT GAGTTGGTGA ATTGTCTTAT
3481 AAGCCATGGA ACTAGTCTCT TAGTCCTGAG AATAGGCCAG TCTAGTTAAA TAGTTATTAG
3541 TTGTGTCTAA TTTTAGGCAG TGTGTTGCAG ATGGGCTTCC ACCAAAGCCA GGCCTCTATA
3601 TGATATGAGT AATCAGTTAT TTAGTAAGAG GCATTTTTGT CTCAAAAAAT AAATAAATAA
3661 AAATATATGA ATAAATGAAT GTATGTTTCT TATCAGACTA CGTCTGTTCT ATCATTAAAT
3721 CCAGAAGGGA GGAGGGTCTG GTTCCCCCTT CCCATCATGG CCTGACCTAG TTTTCAGGTT
3781 AATTTTAGAA CACCCTTGGC TGTGAGGAGT GGTCCATTCT GATGGTTAGG GAGCTTTAGG
3841 ATTTTACTTT TGGTTTACAA AGTAATGTGA ATTAAACAGA CATTTGAGTT AAAGTTTTTA
```

FIG. 4<sub>CONT'D</sub>

4 / 20

3901 TTTTAAATA AAATATTGA TTTAAGCATT TTTTAACTG AATTAATTAG AGCTCTTTTA  
3961 TATATTTTGA TAATGGAACA TTACATACAC AGGCACATAT AAATATATAG ACACATAAAC  
4021 AGAAGTAGAG CTTATAGATT TATACTTTTT TTTTTTTTTT TTTTTTTAAT GAGACAGGTT  
4081 CTCCTTCTGT CATCTAGGCT GGAGTGCAGT GGTGCCATCA CAGCTCACTG CAGCCTTGAC  
4141 CTCCAAGGCT CAAGCAATCC TTCTACCTGA CTGGCTAGCT GGGACTACAG GCGCGTGCCA  
4201 CCATGCCTGG CTAATTCGTG TATTTTTTGT AGATATGGGG AGTTTACCA TCTTGCCCAG  
4261 GCTGGTCTTG AACTCCTGGG CTCAAGAAAT TTTCTAACT TGACCTCCCA AAGTGTTGGA  
4321 ATTACAGGCA TGAGGCACTA CGCCAGACCA GATTTTTTAT TTGTCAGTTT CTAGGTAGTT  
4381 TTCCCCAACT TCAGACTATC AATTTTTTAA TTATCTGTTT TATGCTTAA TTATTAACTA  
4441 GGCAACTCTA AACTTGTATC TCTAAGACAT GACTTTTAGA TGAAATAAGG TAGAAAATGT  
4501 ATATTTCAAA GGCATAGAAT TTAGATCTAA ATAAAGGTAA AGTTATCTAA ATTTTAAGCC  
4561 ATTGTCTTTT CTATTCTAAA AGGTTTTTGA GGTGTTGGTG TAGAGAGGGA GATGCCTTTA  
4621 CAAATGGAAT TTTTGTGTGTT GTTTTTGTGTT TGAGACGGAG TCTTGCTCTG TCACCCAGAG  
4681 TCTCGCTCTG TCGCCAGGC TGGAGTGCAG TGGCACGATC TCCGCTCACT GCAACCTCTG  
4741 CCTCCCGGCT TCAAGTGATT CTCCCACCTC AACCTCCTGA GTAGTGGGA TTACAGCTGT  
4801 GTGCCACCAC GCCCAGCTAA TTTTGTATT TTTAGTAGAG ACCGAGTTT ACCATGCTGG  
4861 CCAGGCTGAT CTCGAACCTC CACCTCAGGT GATCCGCTCG CCTTGGCCTC CCAAAGTGCT  
4921 GGGATAACAG GCATGAGCCA CTGCACCTGG CCTTTTCTGA GTTTTTTAAG GAGTCTGAGT  
4981 CATTAGAAGT CTTTCTAGA TTTTTTAAAA ATGTGGTATT GAAGATGGCA AAGAGGAAGG  
5041 AGGAATAGGG TGGAGTAAAA GTAAATGGGA GGATAGTTT TAAGAAAGGA AGTGAATAGA  
5101 GACATCAAAC ACATTTTAAA AAAAAAGATT TAGTCTACTG AACAAAATTT TTTAAATAG  
5161 GATTTAAAGA GAAAACACAG AAGGCTTTAA AAATATACAC ATAGCTTGAA TATTAGCTTT  
5221 TAATTAAGCT GACTTCTAAC CATGGAGCTC TTTAACAAAA ATTCTTTTAA ATTTGTCTCT  
5281 CTCCTCCTTT AAAACTTTTT GTAGAGATGG GGTTCGCCC TGTTACCCAG GCTGGTCTCA  
5341 AGTCCGGGCA ACTTCTGGGC TAAAGTGATC TGCTGTCTC GGCTCCCAA GTGATAGGAT  
5401 TACAGGTGTG AGCCACTGCG ACTCACCTTA AATCTCTTGT TACCAGATTT TAGTTGGGAC  
5461 AAATGCTGAT ATTTTAAAAG TCACATAAAT ATTAAGCCGA AAAGGACTGA TTTCTGATTA  
5521 GGAAGGAAAC CTAAGCCACG GTGGGAATTT TAATTATTAA ACTGTAAAAT GGAGCAGCCT  
5581 CCATTGTTAA TTTTGTATGG AATCCAAAGT GGCAGTTTGA GTGTAATTGT TTTAGGTCAG  
5641 GTTTTTGTGC TTTAATTTAA TCAAGACAAT TGTTAAGGAT AGCTGTGACA CTATTATGTG  
5701 TCCTTTTAAT TTGATCTATC AATTCCTTAG AACAAGTAAT TTTTTTAAAT TTAGGAATTT  
5761 TAGTCTAAAG GATTTATCTT TTGGCCATTG ACAATTAGAA TTTTAAATGG GGTATTTAAT  
5821 TCCAATAGCA ACTTAATCCA AAGTTTTCTT TATGTCAAAG AAAACAGAAG CCCAGGAGGG  
5881 ATGAGACCTT GTAAGACAAA ACTCCCCTAG GAGCTTGGA TGTGAAAA TACATGTGTT  
5941 GGGCTCCCAA TCTTTTCATA CTGGCTGTGA TGTTACCTGA AAAATCACAT CCTTGGGATG  
6001 GTGGAGACCA AGCGGGAATA TCCCCATCTA GTCACGTCAT GCTCTCAAGG ACATGAGACA  
6061 AGAGGGAAAC CTCTCACCTT GTTTTTATTT CAGGGACTGG CAGCAAAGTT TGTCATAACA  
6121 GAAGTCAGCA TAACCAGAAC CACGAACTG ACCAGTTTGC AGGGCCAGTT CAAACAGTGG

FIG. 4<sub>CONT'D</sub>



5 / 20

6181 GTTGCAGGCC TGTTCTACCC TAGGGTACCC CTCCTTATGA CAGAACACCA AAAGACAAGA  
6241 CAAAAACGAA GGAAAACGGC AACAAACAAA AAGCTATTTT TGAAAGGAAA ATGGCAACAA  
6301 CAACAACAAA AGCTATTTCT GAAGGGAATG GGGTCAAAC ATGAATACTT ATACCACAAA  
6361 GTACTAAAAA ATATATCAGA CTCACTATAC CAAGGTTAGT CACACACAAA ACCTGTTCTC  
6421 TCATTAATCT TACATTTGGA AAGGAAAAGG GAAACAATGA TTTTACTGT CCACTCATCC  
6481 AGAGTCCACA GAGAGAGGAA AACTGGAAAA CTGGGAGTCT GGCAGGAAAT TCTCACTCCT  
6541 CTGCTGGCTT GCCAGGTTCC TGTATTTCTT TCTCTGTGGC TTCCAGAAAA GCACAATAGC  
6601 TTTGGTGGTC TTATTTGTGA TGCCAACTG TGGTCTTGGC CCCCTAAAGT TTCAGTGAAA  
6661 ATCACTGACA TGAAGCAGAT TAATAGGGAA AAAGGCATAC AAATTTATTA AATACGAATG  
6721 GGAGCCTTTA GAATGAAGCC TTGAAGCTAT AGGGGAAATT GTCTATTTTT ATGTTTAGGT  
6781 TTAACAAAGT ATGGACAGCT GTGTAGAAAT ATGACTGGAC AGAAAGGGCA CGATCTAATG  
6841 TTAACAGACT GAGTGGGGAA ACCCAGCAAG GCCTGTCTGT TGAGATTCCT CCTAGCCTCT  
6901 CTCATTCCCT CTTCTGGTG TGGGGCAGGA CCCTCTCTGG AATGGAGGTT TTATGACCTA  
6961 AGTCAAATAA CGTAGGTCAG ATTTTTTTTT TTTTTTTTTT TTTTGTAGC TGGAGTCTCT  
7021 CTGTCAACAG GCTGGAGTGC AGTGGCGTGA CCTTGGCTCA CTGAAACCTC CGCCCCCTGG  
7081 GTTCAAGCCA TTCTCCTGCC TTAGCCTCCT GAGTAGCTGG GATTACAGGG GTGTGCCACC  
7141 ACGCCCAGCT AATTTTGTGA TTTTATAGTAC AGACAGGGTT TCACCTTGTT GGTCAGGCTG  
7201 GTCTCAAATT CCTGACCTTG TGATCCACCT GCCTCGGCCT CCCAAAGTGC TAGGATTACA  
7261 GGGGTGAGCC ACTGTGCCCG GCCTTTTTTT TTTTTTTTTT TTTTATAGGAA GTTGTATTTT  
7321 GGGCTTTTTA ACTAGCTTGT TTTTAAATTA GATTATTGCC TTTAGGGTGG AGCCCTTTAA  
7381 TAAAAAGGGG GAAGAAAACA TAGGTTTAG GGCCTCATAT TTAAATGGGT AAAGCAGGCA  
7441 TAGCTGGAAG GCAGAATACA GAACCCCCCT AATCAAGGAT CTCATTTTTA TATTGAATCC  
7501 TAGGCCCCCC AAAAGAGGGA AATGTCATGG GACGAGATGT GTGGCATTTT TATCGAGTGC  
7561 CCCACTGTAA AGATGCTCCC CCAAGGCTGG CAGGCAGCCC AGTGCCGATT AGCCACTCT  
7621 GTGCTTAGTC TTTTTTTTTT TTTTTTTTTT GAGGTGGAGT CTGCTCTGT TGCCCAGGCT  
7681 GGAGTGCAAT GCGTGATCT CGGCTCAATG CAATCTCTGT CTCGTGGGT CAAGCGATTC  
7741 TCCTGCCTCA GCCTCCCAAG TAGCTGAGAT TACAGGCACC AGCCACTATG CTCAGCTAAT  
7801 TTTTGTATT TTTAGTAGAG ATGGGGTTTC AACATGTTGG CCAGGCTGGT CTCGAACCTC  
7861 TGACCCCAAG TGATCCGCCC GCCTCGGCCT CCCAAAGTGC TGGGATTACA GCGTGAGCC  
7921 ACCATGCCTG GCGTGCTTAG CCTATTTTTA ATGGGAGTTT CATCCTCAAT GGTGAGTGCT  
7981 TTCATTGTCT TTAGGTGCCC CAGACCATGT TTTTAAAAAT TTAAATGCAC GAAGAAATAA  
8041 GTAGCCCTGT ATAGTAGTAA TACTTTGTTG TGAATAACTG TCATAAGTCA TCTCTAAAC  
8101 TGTATTTTTT ATCTAGTTAT TATATATGAC TAGCTATATG TCTAGTTTTT TAAATAATAC  
8161 AAAGTAATTT ATTTTGGCA TCCTCAAAA CCAAAGAGAT TAGGTAATGT AGTGTAGAAG  
8221 AGAGCAGAGC TTTAGACCTG AGAAGAATCT GCCCATGACT CGTGAACTC CACAACGAAA  
8281 GTAGGAGACC CCAAAAAGG GGTGAGTGTC ATCTTTCTG AATTTTTTTT TTTTTTTAGA  
8341 TGGAGTCTTG CTCTGCCACC AGGCTGGAGT GCAGTGGTGC AATCTCGGCT CAGCCTCCCG  
8401 AGTAGCTAGG ATTACAGGCA CGCGCCACCA TGACCAGCTA ATTTTGTAT TTTTAGTAGA  
8461 GACAGCGTTT CACCATGTTG GCCAGGATGG TCTCGGTCTC TTGACCTCGT GATCCGCCCG

FIG. 4<sub>CONT'D</sub>

6 / 20

```
8521 CCTCGGCCCTC CCAAAGTGCT GGGATTACAA GCGTGAGCCA CTGCACTCGG CCGGTCAGAT
8581 AATTTTTTTTG GCCAGTTTTT ACATAGAGTA ATTTTAGGTT TTATGGCTGG CTTTGGGGCA
8641 AAGGGGTTCT GGTTTTTATA GCTGGTCTTG GGGGAGAATG GAACCGAGTG ACAAGAGGAC
8701 AAGAGAGGGT CAGAGAAAAA CTTCTGCTTC TGAGGCGGCT ATTGAGGCCT TCATTTTGGA
8761 GTATTGTCCT CTAAGCCCCA GCAGTGTCAA ACTGTACACA AACCATACAC AGCAGCCAGC
8821 TCGGGTGCTG TTAGGAAATG GTCTCACTGC TGGGTCTGTG GGGTATGTGT GTGTCTGGGT
8881 GTGTGGCTAC TGTCTGCATC CTCCTCCCCC CTACAGCCTC CCCGCCCTCC CTCCAGCCAC
8941 CCTGGGATTG GTGACTCTCA GCCCCCTCCC TCAGCTCCCC TAGACCCTCC CAGAGCCTTT
9001 ATCAGGGAGC TGGGACTGAG TGACTGCAGC CTTCTTAGAT CCCCTCCACT CGGTTTCTCT
9061 CTTTGCAGGA GCACCGGCAG CACCAGTGTG TGAGGAGAGC AGGCAGCGGT CCTAGCCAGT
9121 TCCTTGATCC TGCCAGACCA CCCAGCCCCC GGCACAGAGC TGCTCCACAG GTAGGCAAGT
9181 GGGAGAATGC TGGATGGACC AGAGCTGGCA CCAGGGGGCT GTTATCTCCT GACTGCCCTT
9241 CTTCTTCCTT TTCTTTCATC TGTGTATTGT CAGGCAGCTA CTAATTGTCA ACCCAGAAGC
9301 TGCTGGGTTT AGACCAGGGT CTCAATAAAT CACACCCCCA CAGAAGCCTG CGGGCACTGG
9361 GCACTGATTC CCCCAGTGT TCTGAGTATT CCAGTTTGCC ACTGCCCTGA CTGTAACATA
9421 TGCTAGTATC CATTCTCATT TTTTAAATTT TTATTTATTT ATTTATTTAT TTTTGTAGAC
9481 AGAGTTTCAC TCTTGTCAAC CAGGCTGGAG TACAATGGCG CGATCTCAGC TCACTGCAAC
9541 CTCCGCCTCC CAGGTTC AAG TGATTATCCT GCCTCAGCCT CCTGAGCTGG GATTACAGGC
9601 ATGCGCCACC ATGCCCAGCT AATTTTTGT TTTTATAGTAG AGACAGAGTT TCACCATGTT
9661 GGCCAGGCTG GTCTTGAACCT CCTGACCTCA AGTGACCCGC CCATCTCGGC CTCCCAAAGT
9721 GCTAGGATTA CAGGTGTGAG CCACTGCGCC CAGCCTATTT CTTTTTTGAG ATGGAATCTT
9781 GCTCTCTCGC CCAGGCTGGA ATGCAGCAAG CATGATCTCG GCTCACTGCA ACCTCCATCT
9841 CCCGGGCTCA AGCCATCCTT CAGCCTCGGC CTCCCCAGTA GCTGAGACCA CAGGCACATG
9901 CCACCACGCC TGGCTAATTT TTTATATTTT TGGTAAAGAT GTGGTTTCAC CATGTTGCCC
9961 AGGCTGGTCT CAAACTCCTG AGCTCAAGTG ATTCACTCGC CTTGGCCTCC CAAAGTGCTA
10021 GGATTACAGG TGTGAGCCAC TGCACCCGGC CTTACCCATT ATCTTTTGAA CATCTACTAT
10081 GCATTAAGCT CTTTACATGC ATTAACCTCTA ATACTTTCAA TAACCCTGTG AGGTAGGCTC
10141 TTTTCTTTCT CCCATTTTGT AGTTAAAAAG CCAAGGCTCA GAGAGGTAA ATAACCTGCC
10201 GGGGGTTCCA CAGCTGTAAG TGGTAAAGCT GGGTTACAAA CTATTTGACT CTAGAGCTTT
10261 TAACCACTGC CTAAGACTGC CCCTCATCAA TAGAGGCTTG GGCAACCCAT GGCCCTAGGC
10321 AGACCTGGGG GCAGGAGGGC TGCATAGGAA AGGGCAGAAC TTTCTAGTTC TAGAACAAC
10381 AATAAAAAGA AGAAAGCCTT CAGAGGCTCC ACATTAATTG GAACAAAGGG GATTATGACA
10441 GATGCTTAGG CATGTTTGTT GAATTATTAA TAAATAAAAT CAGACTAGGG ACTGGGGACT
10501 CCAGTCTTGG AGGCCTTCAC AGGCCCAGAT CCCAAACCCA CCAAACCCAC TAGACCTGCA
10561 GTGGAAGCTA CAATGAGCTT GGATAGTTCC TGCAGTTAAC AGCAATATAC TATGTATTCT
10621 GCCTCTTTCT ATTTAAATTT TTTAACCTGA TATCTTAGTA AAACTTTTTC ATAAAAATTC
10681 CAGACATTTG GAAGTGCCAA AAATCAAGTC ATTTTTTATA TCTTCAGTAA TTCTGTGCCA
10741 TAAACAAACA GGTTGCTAGG TGCTCTATGG GATGTAAAC CTTGGCCAGG CAAGGTGACT
```

FIG. 4CONT'D

7 / 20

10801 CACTCCTGTA ATCCTAGCAC TTTGGGAGGC TGAGGCGGGA ATATTGCTTG AGCCCAGGAA  
10861 TTTGTGACCA GTCTGGGCAA CATAGTGAGA CCTAGACTCT AAAAAAAAAA TTTAAAAATT  
10921 AGGTGGGTGT GGTGGGTCAT ACCTGTAGTC CCAGCTACTT GGAAGGCTGA GGTGGGAGGA  
10981 TCGCTTGAGC CCAGGAGGCG GGCAAGGCTG CAGTGAGCTG TGATGGTGGC ACTGCACTCC  
11041 AGCCTGGGCG ACAGAGCAAA ACCCTGTCTC AAAAAAAGAG GCAAAAACAA AAACCTTAAGA  
11101 ATCCTTGTTT TAGATTGGGG CAGACTAAAG AGTCAGTTGC CATGGATGAA GCTTGATTGG  
11161 ATCCTGGAAA AGGAAAAATA AAGCTTCAAA GGACATGTTT AGAAGTTTAT AAAGGACATG  
11221 TAGAGAAATC TGAGAGTGGA TCGCTGTTGG ATGAGTGATG TTGATTTTCT TAGGTGTGGT  
11281 GATGGAGTTA TGATTGTGTA AGAGAATGTT CCAGTTCTTG GGAGAGGCAT GCTGACATTT  
11341 TAGGGTAAAA TGTCATGATA TCTATAACCT ACTTTAGGAT GGTAGGGTAG CAAGGATTTG  
11401 TGTAAATGTG TATATGCATG TATTTATATG CACACATATG TGTGTGTGTC AGAGCACACA  
11461 GATAGTGCAA GGTGTTAACA TTATCAGTTG GTGCATTTAG ATGAGGAACA TACAGTATAC  
11521 AGATGTTAAT TGTATCTTTT TTCAACTTTT CTGTAAGTTA AAAAACTTT CAAAATAATA  
11581 AGCTATATTG AATTTTAAA ACATCATATT ATGCTATTCT TCTGTATAAA TTCTCCAATG  
11641 GTGTTCCATT TCACTCCTTA CCACAGCCTA CAAGGCCCAT CATGATCTGC CCCGACCTAC  
11701 TCTCTGATCC TCTCTCTTCC TGCTCAAGTG ATTCTGGCCA CCCTTTTTTT TTCTTCTTTT  
11761 TTAGACAGTC TTGCTCTGTC ACCCAAGCTG GAGTGCAGTG GTGCGATCTT GGCTCACTGC  
11821 AACCTCCACC TCCCGGGTTC AAGCGATTCT CCTGTCTCAA CCTCTAGAGT AGCTGGGATT  
11881 ACAGGCATGC GCCACCATGC CCAGCTAATT TTTGCTCACC CTGGCTTTTT AATGTCTCTG  
11941 GAATATGCTG CCACTCATTC CTGCCTCAGG GTCTACTTCT TTGCATCACA GCAGATGCCA  
12001 TTATCTGACA TCACACTATA TATTTATTG CTTGTGTAGT TGGTCCCTT CTCCACCCTA  
12061 CAGTAGAATG TAAGTCCAGT GAAAATGAAG ACTTTGTTCA CTGTTATGTC CCAGTACCTA  
12121 GAACAGTTCC AGGCACTAAG TAGACACTCA ATAAATGTTG ACTAGTGAAA AAAAATGTGA  
12181 GACCTGGGAT CCTGCCTTAT AAGGACTCAG TGTCTAGAAA AGGGAGCTGT TTTCCATGCA  
12241 AATAACTGTA GTACAAAGAC GAGTGTAGGC AAATTGCTAT GGGGCTTCAA AGAAAGGAGA  
12301 GGCAATCCGG GGCTTGGGGA ATCAGGGAGG GCTTTGAGCT GATCTCCAG GTTGGCAGAG  
12361 TTGAGTCAAG AGAGCATCGA GAGCTAAGGC ACACAGTGAT CATGCATGGG CTGGGTAGGG  
12421 GCATGGGAAA GAGTCCTGTC CGGGTGGTGT GCCCAGGGAA TGCAGGGGTC CTGCGACATG  
12481 AGGCTGGGCT CTTAAGTGTC AGGGAGGAAA CCCAGGAGAG AAAAGCACTT CCAGTGAAAC  
12541 CCTGGGAAAG GCCAGAGAGA AGGAGGAAGA GCATGGGATC TTGGACAGAG GCTGGAGCAA  
12601 ATTGTAAGTG ACCTCCGCTG ATTGGATTTT TGACCGTGGT TAGGACCCTG ACTATTGCTC  
12661 ATTCAGACAT GAGACACATT TGCTTACAG CTCTCTTTGT TGTTCGAGGG TCTGGATCCC  
12721 TCAGCTTAAG AGAGGAATGG GGGCTCTGAA GCTCTGGGCC TCTTCATTGT CTCCCTGAAT  
12781 TCATTTGCTC TTTCTCCTTT GCTCCTTTAT TTGCTCCTTC TTTCTTTGAA TGGAGGCTGA  
12841 CATGTTTGGA CTTGACTGAT TTGAGAGGAG GGGAAATTTG GTACCTAGCC AACAGCTGAC  
12901 ACAGACAGTG GCTGCCACCT GTAGGCAATT GTGAACAGAA GGAATAGAAA GCTACAGGAG  
12961 CAAAACCTTG AGACCAGCTT TCATATTGGT TCCTCTTACC TCACTGCCCT GGGTAGCAGG  
13021 TCTTTGGTTG GAACTAATCG TTCTCTCCCT CCAGTCTCCT ATTCATGCTC TTACCTCCCG  
13081 GCCTCAAGCC TGCACCTCTT GCTGAAAAAG ATCCAAGAGG TGACTCCCTT CCATCTCTTC

FIG. 4<sub>CONT'D</sub>

8 / 20

13141 AGCTCCACCC CTTGCTTCTC ACTGTGGGTT AACTTCCTCC TTTGAAGTGG CAGGATCTGG  
13201 GTGCCAGTTT GCCTGTCAGG AAGTGTCTTCT TATCACTCCA CTCCCAATCC CCCTGGTCCC  
13261 AAACCTAGGTA CAGAAATTCC TACTGGGGCT GAAGAACAAT TTGCCATCCA CAAACGTCTT  
13321 AGACAAGACA TGGCCAGCCG CCCCTACAA GTGCCTCAGC ACAGCAAATC AGGAGCTGCA  
13381 GCAGCTCTTC TACCAGTGGA AGGCAAGTGG AGCCCAGGCA CCCCTCCTCT CATTTCTGCT  
13441 TTTTTTTCCC TCCCCCTGAT TTTCTCTTT TGCCCTCCCTC TTCTATTTTT TTCCCATTAA  
13501 AAAAATTGTG GTAAAATATA CATAACATAC AATCTACCAT TTAAACGGTG TTTAAGTGTA  
13561 TAGTTCAGTG GCATGAGCGA CATTTCATGTT GTTCTGCAGC CATCACTGCC ATCCATCTCC  
13621 ATATGCGTTT TTCATCACCC CAAACTGAAA CTCTGTACCC ATTAAGCAAT AACCCTCTAT  
13681 TCTCCCATTC CCCTAGCCCC TGATATCTTA TAATCTACTT TCTGTTTCTA TGAATTTTAC  
13741 TTTTCCAAGT GCCTCATATA AGTGGGAATC ATATTTGTCC TTTTGTGTCT GGCTTATTTT  
13801 ACTTAGCATA AAGTAATTTG TTCTTTTATT CAGGAAATGC TTATTGAGCA CCTGTCTGGG  
13861 ACTAAGCCTT GCCCTGAGAG CTGAGCATAG AGCCCTCCTG GTGCTTTTAT TTGATGGTGT  
13921 CCATTCCCTC CCCTAGCCTC CCTCAGTTCT CGCACTCCTC CTCAATGGTC CTCCAGCCCC  
13981 GGCCTCTCCC TGAGGTGTCT AGTGCCTGTC CTTTTTCCTC AGTCTCTCTC CTCTCCTAGT  
14041 GTCTTCTAGT CAATATTTCT CACCTCCCTC CCCAGCCCTG CCCTCCCACT CTATGATTTT  
14101 AGCTCCTGTC CCTCCTTCCT CACAGTGCAA GAGGTTCCGG GATCAGCTGT CCCCAGGCA  
14161 GGTAGAGATC CTGAGGGAAA AGCTCTGTGC CAGTGAAGT TCAAGGGCA AGAAGGCTTC  
14221 ATATCCCCAG AGGTGAGGGC CTCCCAGACC CTGCACAGCC AGTTCCATCA CGCAGCAGTT  
14281 CTCAAAGTTC AGCGTGCCTT AGAATCACCT GGCAGGATTG TCACCCCCAG GTGCTGTGTC  
14341 CCTCCTCAGA GTCTCTGATC CAGCAGGTCT TGGGGTGAGG ACCAAAATTT GCCTTTCTAA  
14401 CAACTCCCCA GGTGGTGCTG ATGTCTTGGT CCTGGACTGT GCTCTGTGGA CACTGACAGA  
14461 GGATACGTGG ATGTGGGGGA AGGGCCCCGG AGGACTAGGA TGGGAACTCT GGGGGTGGGG  
14521 AAGAGGCCTC TGGGCCCTGT CGCGCTGCAC ACCTCCCATG TGTCTCTAGT GTCCCCATTC  
14581 CATTCTGTGG TGAATACATT GGGCTGCAAG GGAACCCCAA GCTGCAGAAG CTGAAAGGCG  
14641 GGGAGGAGGG GCCTGTTCTG ATGGCAGAGG CCGTGAAGAA GGTCAATCGT GGCAATGGCA  
14701 AGGTAAGGGC CTGCAGGCTG AACTCCTCCC GCAGCTAGTG CAGAGCTGTG GGCTGGCATC  
14761 TGGAGAGCAG ATGGCAGGCT GTGTTTGCGC CCTGCCAGGT GGAGTGGGGG CAATTAATCC  
14821 TGCCCTTTCCT CACCCTTGCC TGTTCCGTCC CTAGACTTCT TCTCGGATTC TCCTCCTGAC  
14881 CAAGGGCCAT GTGATTCTCA CAGACACCAA GAAGTCCCAG GCCAAAATTG TCATTGGGCT  
14941 AGACAATGTG GCTGGGGTGT CAGTCACCAG CCTCAAGGAT GGGCTCTTTA GCTTGCATCT  
15001 GAGTGAGGTA TCAGAGCTGG GTGGGGCAAG CCTTGGACTG GAGAAGGTGG TATGCATCCC  
15061 AGGGCTGGGG CAGGCTGGAG GTGATGGGGA CCAGACCTTT CGCTCTGGGC CTTTGATGTC  
15121 CCTCAGGTGC TCCTGAAGAG AAAAAATGAA TCCCTTTCCT GCTATTTTTT CCTCTTCCTA  
15181 AGATGTCATC GGTGGGCTCC AAGGGGGACT TCCTGCTGGT CAGCGAGCAT GTGATTGAAC  
15241 TGCTGACCAA AATGTACCGG GCTGTGCTGG ATGCCACGCA GAGGCAGCTT ACAGTCACCG  
15301 TGAAGTGAAG GTGAGGCCAT GAACTGGGGG TGAGGGGCGG CTTACGGTAG ATGGCCAGGC  
15361 TGATGGTCAT CGTGACCAGG ATCAGAAAGC GAAGCATGTA GGGCAGTGCA GGCCGGGGCT

FIG. 4<sub>CONT'D</sub>

9 / 20

15421 TGGAGGTGTT TCTCAGGCC CCACCCAGGT TCTCTGGGGC CTCAAGTCCT CTGACTCGCA  
15481 TGATGGGGGG GCCATCATGG AAATGCGGGA GTCGGGGTGA GGGGATGGGC ACTAGACTTG  
15541 GTTTTCTGTT CCGTCTCCAG GTTCTCAGTG AGGTTCAAGG AGAACAGTGT GGCTGTCAAG  
15601 GTCGTCCAGG GCCCTGCAGG TGGTGACAAC AGCAAGCTAC GCTACAAAA AAAGGGGAGT  
15661 CATTGCTTGG AGGTGACTGT GCAGTGAGGA GGGGGCACCA TGCAGAGATG GCAGTTGCTT  
15721 CCTCCTGAAC CAGCACTAAT CCCCCTCTGC CCTCCTGTGT GGGAGGATCT CTAACCCCTC  
15781 TGATCGTGGC GCATGGCTTG GGGATTAAAC TACCCTTGAA GAGGACCCTT GTCCCAAACC  
15841 CTTCTTGTTT TCTCCTCCAA AAGTAGCTTC CTCCAACCCG CAGCCTCTCT GCACACTAAT  
15901 AAAACATGTG GCTTGGAAG GTTCAGTCAG GGTGGGTGGG TCCTTGTTCC CCCTATCTTT  
15961 TCACCCAGGT GTACTTAGAC CCCTGCCCC ATGCCCTTTT TCCTCCTCAA GCTCCTTGGA  
16021 GCCAGCTAGT GAGGTAATAA GAAAGGAAAA GAAGGAAAAT TGTCTCCGGG CTCCTTGACC  
16081 GGCTGAGCTC TGGGGGGGTG TTTAGAGAGA CTGCGGTGGG TGGAGGGGCT GCGGGGGGAG  
16141 TTAAGGATGG GGCTCAGGTC GCAGGTGGCC AGTGGACTGA TTCATTAAGT GTGTCCCTGG  
16201 AGGAAAGAAG TGAGCATCCC TGTCTTGCCA GAAACTGGGG TCCTTTGGCG ATTTAGCCTG  
16261 AAAAGCAGCC CAAGGCTGGA GGGCTTATGT ATGCTGGGGT GCTGGGGAAT GCAGGGTCTC  
16321 CTGTACTTGG GAACGCCATC ACCCCTTCTA CTCCCACACA CAGCACAGGG CTCCATCACA  
16381 CCAGCCTCCC CGACACCCCC TTCCTTCTCA CACACCCGAG ATGCCAAACT GCTGCCAACA  
16441 GTTATCTTGC TCGTCTCTGT CCCACAGCTG GGGCCTGCAG CAGGTGGCAC TTCACATCAC  
16501 TCACTTGATG AGGCTCCCTC ATCAAGACCC TCCCATCCCT GTAACCTGGC CTTTCCCTCT  
16561 CCTCTTCCTT TATTTTTCCT GCGTCATTGT CATTATCTTT TTCTCACCCT CCAACTATC  
16621 TCACACCATC TCATTGTCCC TGTTTCTGTG AGCTCTGACT AATATCAATA TGTAATATTT  
16681 TGTAATAATG TTTAAATATT TTCCTACTCC CCCTCATATC TATTTTCTCA TAGATTCTGT  
16741 CTTGTCTGTC TTGTCTCTAC CTTCTGTCTG GCCTCTACCT TTGGGGAACA AGCTGCTCAT  
16801 GTAGTCACAG TAAATTTAG ATCTGTGGTC TGTGAGAGCT TAGCAGGGTC TGCCTTTGTT  
16861 TTTGTCTCTG GCTGTCTCTT CCTCTTCTCA AGATCTCTAC CTTGCCTACC TCTTCCGCT  
16921 TCCTTCCCTT AACTCACTAT GCCTTGGGGC TGGGGTCTCC CTCCACCTGA CTTCCATCTG  
16981 CAGGCAGCTC ACGGCCGGCT ATCATGCTGG CCAGGGAGAA CTGATTAAC TCTCTTCCTG  
17041 CCTGCAGATT AATCTGCTGT CTGAGCACAA GCCACGTGCT TCTGGCACAC CCTGCTTTGA  
17101 GCTGAGATAG AACCTGGGGA ATCATCTGTT TTCAGGCGGG TGAGGGGCTA GAGCCTGCCT  
17161 TGTTTGGGAG GAGGGTGGCT CTGTTCAGAA TAGGGGTAGC TCAGGCTCTG GCCAGCCTTC  
17221 TCCCGCCCC AACAGCTCCC CCCATCCTTG ACTTCTCAGA ATCAGGCCGA GAAGAGCCTA  
17281 TCTGGCCGAG AGTGGGGTGG TGACCTGCGC CTCATCGCCC CCGCTCTCCA TCTCATCTCC  
17341 TGCTCCCAGG GCCCAAATTG TCGTCACTTT CCCAGTGAAG TGTCTGGTCA TTTTCAGAAG  
17401 CAATTTAGG AGAACATGCA GCTGCCGCTC CCTATCCTGC ATTTCCCTTC ACAGGGCTGA  
17461 AGGCACTGTC AGCTCCCTGG GCTGGGGGTG ATGGGAGAGG GGAAGGGCTA GGGCCCTCAC  
17521 CCCTGTCCCT ACTGTGCCCC TCATGTAGAT GGACTGGAGT TCAAGGAAGG GCAGGCACTC  
17581 CCCTCCTCCT TTA CTCTCTCT GTCACTCTCT TCCTCCTCTT CTTTCTGTG TCTGCCTCTC  
17641 TTTTCTGGAG CCTAGGAGTG TGTGTTTTCA TCCCCTGAAA CAAATAGGGA CTCAGTTTCC  
17701 CCACCTGTGT TACAGGGTTG GAATTGGCTC CATCACTGTG GGAGAAGCTG GAGTTCTGTG

FIG. 4<sub>CONT'D</sub>

10 / 20

```

17761 ACCAGTCCTC CCCTCCCCAG CCCTGCCTCT TCTCTCCCAG CCCTCTCCCT TCAGCCAGTT
17821 CAGCGCTCTG AGAGTCTGGG TTGTTTCAGC CTCTGAGGGG CACAAGCCAT CCTGGATTCC
17881 CCTAACCCCA TGAGGAGCCA TTCTAGCATC TCACAGCTTA AACCAGCTCT AGCTCAGTCC
17941 TCCTGGCTTA GTCCATTTTT CTTCCTCAGG CTCTGAGGGC CTCTTGTTC TTGCTCTGTG
18001 GGGTTTTCTC CAGTTGTCTC CTGGCTGCAG GACATGGCAG GACATAGAAT GCTGTCATCC
18061 TTCCACTCTT CATTGGCATC TCCACCCAGT GTCACATATG ACCCTAGCCC TGCTCTCCCC
18121 TTGCCAGTAC CCCTCTGGGA TTTTGCGAGA GTCCACAAGT TGTGCATGTG GTGGATATAT
18181 TCAGGCCATC TTGTGTGTAC AAGCTAGAGG GTCTGCTTCC ACCCTCTGGCC CTCAGTGAAT
18241 TGCTGACTAA CCTGTCTCAA CACAGCACAA CTGTACACAC CTTTTCTGG CCTCATCCCT
18301 AACCACATCAT AGCAGCAAAG AGGGGAAGTT GCAGGGGAGG AGCTGCTAAG GACCCTGGAC
18361 TCCAAGTACC CTGCTCCTCT AGGCCAGGGA CATCATCTGA GATGTGGCTC AAATAAAGGG
18421 TGGGTGTTCA AGAAAAACA CTTGGGGACT CTATAGCTGC AACACCCACT TTACATGTCA
18481 TTTCCATATG ATTTGTAGGC AAAATGAAGC CCAGGCTGTC CTAGCCCTCC AATACCTCCC
18541 TCTCTCATCA CCTCTCCAAC ATAGCCTAGC ATTAGCTCTT TCAAGTCTTT GCTAATCCCA
18601 GAGATCAAGG GGTGATCAAC TCTCCCTGCC ATCCCCTTGT TCCCCGCACC CCCCCCCCCG
18661 GTCCCCCAC CATCCTTGGC TCCTGCCATC CTCTTTGAGA TGCTGCATCA TCAAAGGACA
18721 TTATTTATGG TGTACCTTTG CTGAAGCCCT GCTTCCCTGG TGCCAGGGCT TGGGAGCAGG
18781 GATGGGTGGG TTGGTGGGGG AGAGGGGTGG ATGCAGAGAT TGGACCCAGG AGGCTTTTAG
18841 TCCTCAGCTC TTGGCTTAAC ACCTCCTCCT CTTACACACC CAACTCCCTC CAGCCTGCCC
18901 AGCTTGGGCC TTCAGCTCCA GATTGGTGGG GTTAGGAGAG GAGGAGGAGG GAGATGGATG
18961 GAACCAATTA GGAACAGCAC CTGGGCTCCT CACAGGAATG AACCAGTCAT GCCATTTGCA
19021 TGTAACACAG TTCCCACTTC TCTCCTCATC CTACCAAATG CTCCCAACCC TGGGTTCTGG
19081 CCCATGTTCT TTGCCCACAC AGCCCTGTAA TTAGCTGGGT AATGAGAAGC TTTTAATGAG
19141 TCCCATTAGC ATCTCGTGTA ATAAAGAGGC CTTGAGACCC AGCTGCTGTC CTCACTTTGG
19201 GATGAACACG GGTCCCTGTG TAGCCAGTGA CTTCTGTCAG TACAGTCTAA GTTCTCGGAT
19261 GGGGTGGGAG ACAAACATTT CAGGACCCCA GCAGCACTTG AGAGGTTCCTA TGGTGGATCC
19321 ATGTTTTTGA CTGTGATACA AGAACTTGG CTCTGGCTTC CTTGTTTATT TTGTAAATAA
19381 CATTTTTTCT TCTTTTAAGA GACAGAGTCT TACTTTGTTG CCCAGGCTGG AGTGTAGCAA
19441 TGCAATTATA GCTCACTGCA GCCTCAACCT CCTGGGCTCA AGTGATCCTC CTGCCTCAGC
19501 CTCTGGGATA GCTGGGGCCA CAGGCATGCC CCACCATGCC TGGCTAATTT TTAAAAATGT
19561 TTTTGTAGAG ATGGGGTCTT ACTTGCTATG TTGCTCAGAC TGGTCTCGAA CTTCTGGCTT
19621 CAAGCAATTC TCCCACCTCG CCCTCCTAAA GTGCTGGGAG TATGGGCATG AGCCACCATG
19681 TCCAGCCTTG TAAATACATT TTTATTGAGC ACCTATTATA TGTCAAACAT TATAAAGTGA
19741 GGGATACAGT AGCAAACAAA ACAGACAAA ATTTTGGCCA TCATGACACT TATATTCCTG
19801 GGTGGGAGTG GTGATAGAAA GACAATAAGT AAAATACTTA GCATAGTGGA TGTAATAAGT
19861 TCATGAAGGG AAAAATGGGA GTGAGGTATA TGGAAATTTG GGGTGGTGAT AATTTTAAAT
19921 AGGGTGATTG GGGAAATGCTT TGTTGCACAG ATTGTTTTTG TAGTAAATAT GAGATAAAGA
19981 TACGGTTCTC TCCCAAACCTC AAAATGTAGA AGAGTAGAAG GTCCCAAATC TTCAAGTCTC

```

FIG. 4<sub>CONT'D</sub>

11 / 20

20041	TTGGAGAGGG	GGGCCACCCA	TTCCGTCTGG	GACAGTTAAC	TGTTCCCTCA	CAGGTCAAAG
20101	TTTATGCCAG	TGCAGTAAAA	AGAGTGGGAG	ACCTGGGGTG	AGACAAACCT	GGATTGAGG
20161	CTGTTCTTCA	CTGATTAGTA	GCCATATGTA	CTGGAGCAAG	TGACTGAACC	TTCTGAGCCT
20221	GTTTTCTCAT	CTGGAAAATC	AGAATATTTT	CTACTTACAT	GGTCATGGTG	ATGAAAACCA
20281	GATGGACTGC	TCCATGCCAA	AGCACCTGTC	AAACATTCAA	ACCCTGCACC	CATTACAAAT
20341	ACTGGGCTGA	CGGATGGCTC	TGGCTTTGCT	TTTGCATCTC	CGCTGTCTCA	TTCAGCAGCA
20401	GCATCTGGCT	CTGGCTCTCG	GCTCTGATCC	TGGTTCTGAC	TCTCCCCTGG	AGCTCTCTCC
20461	CTTGGGTGAG	AAATAAGCAG	ATAATCTCCC	TCATCTGTGT	GTGGTGTGAA	CAAGAGGCTT
20521	GAAAGGTCAG	AGAAGAAGAT	GCCTGAAGTG	CAGGGAGACA	GATTAGAGTG	GGGAAAATGT
20581	AACTCTGAGG	AAAAAGGGAA	GCAATTAAGA	GATCAAGGCC	AGGGGCAGTG	GCTCATGCCT
20641	GTAATCCCAA	CACTTTGGGA	GGCTGAGGCG	GGCAGACCAT	GAGGTCAGGA	GTTCGAGACC
20701	AGTCTGGCCA	ACATAGTGAA	ACCCCGTCTC	TACTAAAAAT	ACAAAAAAT	TAGCCAGGTA
20761	TGGTGGTGTG	CACCTGTAAT	CCCAGCTACT	TGGGAGGCTG	AGGCAGAAGA	ATTGCATGAA
20821	CCCGGGAGGC	AGAGGTTGCG	GTGAGCCGAG	ATTGAACCAT	TGCACTCCAA	CCTGGGCAAC
20881	AGTGTGAGAC	TCTGTCTCCA	AAAAAAAAAA	AAAAAAAAAA	AATCAAGGCC	GGGGAGGGGG
20941	CAGGGGTGGC	ACAGCTATCG	AGTTCTGTTC	ATCCTCTGTG	AGATTACATC	AGGAGGTGTA
21001	AAAGAACTCT	AGAAGAATGA	AGCTAAGTCC	AGCTGATTCA	GGGTTCAAGA	AGGATTGAGG
21061	TGGGAGAGGC	ATCATGACCA	CTGGTGAGGA	GTGGAGGAAG	GCCGACACTG	GAGCTTTCTT
21121	TGCCCAAGCA	GAGGAGGGGT	GTGACACTCT	TGAGGACCAA	TGTAATGGCG	CAGCTCCCTC
21181	TGGGAGGGGG	AAAGGAGAGG	ACTGGAGGGG	ATGCTAAACT	GACCTTCTAA	CCTTCAGGGG
21241	CCTGAGTCTG	GTTGTCTTGG	GTGGGGAGGG	GCGCCTGCCT	GAAACTGTTT	TAGCCCAGAA
21301	GTCAGGCCTG	AAGGTTAAAG	GGCAAGGAGC	TGGTGGATGA	ACAAGGTGGG	GAAAGAGGCC
21361	CAGGGTCCAC	ATCTACTGAG	CTGGACTCAG	GCATGGGAAT	TGGTGTGTG	AGGGCCAAGA
21421	CACCTGGCCT	CCTAAAAGTT	TGCTGAAAAT	CACCTGACATG	AGAGTAATTG	ATTTATAGGA
21481	GAAAAGGTAG	ATAAATTTAT	TTAATATGTA	TATATGAGCA	CCTTTAGAAT	GAAGACCCAA
21541	AGATATAGGG	GAAATTGCCA	GTTATTTATT	TATTTTTTTT	GGAGATGGAG	TCTCACTGTG
21601	TCTGCCAGGC	TAGAGTGCAG	TGGCAATGAT	CTCGGCTCAC	TGCAACCTCC	GCCTGCTGGG
21661	TTCAAGCAAT	TCTCCTGCCT	CATCCTCCTG	AGCAGCTGTG	ACTACAGGCA	CGCACCACCA
21721	TGCCCCGGCTA	ATTTTTTGTA	TTTTTTAGTA	GAGACAGGGT	TTCACCATGC	TGGCCAGGCT
21781	GGTCTGGAAC	TCCTGACCTT	GTGATCCGCC	CGCCTTGGCC	TCCCAGAGTG	CTGGGATTAT
21841	AGGCGTGAGC	CACCGCCCCC	AGCCTGAAAT	CGCCAATTTT	ATGTTTATGT	TTTACAAAGT
21901	ATGGACAGCT	GTGTAGAAAT	ATGACTGGAC	AGAAGGGCAT	GCTCTAATGT	TAACAGACTG
21961	AGTGGGGAAA	CCCAGGAAGG	CCTGTTGAGA	TTCCTCCTGG	CCTCTCTCAT	TCCTTCCTTC
22021	TGGGTATGGG	GCAGGACCCT	CTCTGGAATG	GGGAGATCTT	AGGACCTAAG	TTAAATAAGG
22081	TAGGTCAGAT	AATTTTTTAT	GGCCAGTTTT	TACATACAGT	AATTTTAGGT	TTTATGGCTG
22141	GCTTTGGGGA	AAAGAGGTCC	TGGTTTTTAT	AGCTGGCCTT	GGGGGAGAAT	GGGACCCAGC
22201	AACAGGAGGA	CAGGAGAGGG	TCAGAGAAAA	ACTTCTGCTT	CTGAGGCTGC	TACTGAGGCC
22261	TTCATTTTAG	GGTATTGTCT	TCTGAGCCCC	AGCATTCCTC	GGTGTGAAAA	ATTTTAAAGA
22321	AATTTTATAG	TCCAGAAATT	GAGTTGGTGA	ATTGTCTTAT	AAGCCATGGA	ACTAGTCTCT

FIG. 4<sub>CONT'D</sub>

12 / 20

22381 TAGTCCTGAG AATAGGCCAG TCTAGTTAAA TAGTTATTAG TTGTGTCTAA TTTTAGGCAG  
22441 TGTGTTGCAG ATGGGCTTCC ACCAAAGCCA GGCCTCTATA TGATATGAGT AATCAGTTAT  
22501 TTAGTAAGAG GCATTTTTGT CTCAAAAAAT AAATAAATAA AAATATATGA ATAAATGAAT  
22561 GTATGTTTCT TATCAGACTA CGTCTGTTCT ATCATTAATT CCAGAAGGGA GGAGGGTCTG  
22621 GTTCCCCCTT CCCATCATGG CCTGACCTAG TTTTCAGGTT AATTTTAGAA CACCCTTGGC  
22681 TGTGAGGAGT GGTCCATTCG GATGGTTAGG GAGCTTTAGG ATTTTACTTT TGGTTTACAA  
22741 AGTAATGTGA ATTAAACAGA CATTTGAGTT AAAGTTTTTA TTTTTTAATA AAATATTTGA  
22801 TTTAAGCATT TTTTTAACTG AATTAATTAG AGCTCTTTTA TATATTTTGA TAATGGAACA  
22861 TTACATACAC AGGCACATAT AAATATATAG ACACATAAAC AGAAGTAGAG CTTATAGATT  
22921 TATACTTTTT TTTTTTTTTT TTTTTTTTAA TGAGACAGGT TCTCCTTCTG TCATCTAGGC  
22981 TGGAGTGCAG TGGTGCCATC ACAGCTCACT GCAGCCTTGA CCTCCAAGGC TCAAGCAATC  
23041 CTTCTACCTG ACTGGCTAGC TGGGACTACA GGCGCGTGCC ACCATGCCTG GCTAATTCGT  
23101 GTATTTTTTG TAGATATGGG GAGTTTTACC ATCTTGCCCA GGCTGGTCTT GAACTCCTGG  
23161 GCTCAAGAAA TTTTCCTAAC TTGACCTCCC AAAGTGTTGG AATTACAGGC ATGAGGCACT  
23221 ACGCCAGACC AGATTTTTTA TTTGTCAGTT TCTAGGTAGT TTTCCCCAAC TTCAGACTAT  
23281 CAATTTTTTA ATTATCTGTT TTATGTCTTA ATTATTAAT AGGCAACTCT AAACCTGTAT  
23341 CTCTAAGACA TGACTTTTAG ATGAAATAAG GTAGAAAATG TATATTTCPA AGGCATAGAA  
23401 TTTAGATCTA AATAAAGGTA AAGTTATCTA AATTTTAAGC CATTTGCTTT TCTATTCTAA  
23461 AAGGTTTTGG AGGTTTGGGT GTAGAGAGGG AGATGCCTTT ACAAATGGAA TTTTGTGTGT  
23521 TGTTTTTGTG TTGAGACGGA GTCTTGCTCT GTCACCCAGA GTCTCGCTCT GTCGCCCAGG  
23581 CTGGAGTGCA GTGGCACGAT CTCCGCTCAC TGCAACCTCT GCCTCCCGGC TTCAAGTGAT  
23641 TCTCCACCT CAACCTCCTG AGTAGTGGGG ATTACAGCTG TGTGCCACCA CGCCAGCTA  
23701 ATTTTTGTAT TTTTAGTAGA GACCGAGTTT CACCATGCTG GCCAGGCTGA TCTCGAACTC  
23761 CCAACCTCAG GTGATCCGCT CGCCTTGCCC TCCCAAAGTG CTGGGATAAC AGGCATGAGC  
23821 CACTGCACCT GGCCTTTTCT GAGTTTTTTA AGGAGTCTGA GTCATTAGAA GTCTTTTCTA  
23881 GATTTTTTTA AAATGTGGTA TTGAAGATGG CAAAGAGGAA GGAGGAATAG GGTGGAGTAA  
23941 AAGTAAATGG GAGGATAGTT TTTAAGAAAG GAAGTGAATA GAGACATCAA ACACATTTTA  
24001 AAAAAAGAT TTTAGTCTAC TGAACAAAAT TTTTTAAAAT AGGATTTAAA GAGAAAACAC  
24061 AGAAGGCTTT AAAAAATATAC ACATAGCTTG AATATTAGCT TTTAATTAAG CTGACTTCTA  
24121 ACCATGGAGC TCTTTAACAA AAATTCTTTT AAATTTGTCT CTCTCCTCCT TTAAACTTTT  
24181 TTGTAGAGAT GGGGTTTCGC CCTGTTACCC AGGCTGGTCT CAAGTCCGGG CAACCTCTGG  
24241 GCTAAAGTGA TCTGCCTGTC TCGGCCTCCC AAGTGATAGG ATTACAGGTG TGAGCCACTG  
24301 CGACTCACCT TAAATCTCTT GTTACCAGAT TTTAGTTGGG ACAAATGCTG ATATTTTAAA  
24361 AGTCACATAA ATATTAAGCC GAAAAGGACT GATTTCTGAT TAGGAAGGAA ACCCTAAGCC  
24421 ACGGTGGGAA TTTTAATTAT TAAACTGTAA AATGGAGCAG CCTCCATTGT TAATTTTGTA  
24481 TGGAATCCAA AGTGGCAGTT TGAGTGTAAT TGTTTTAGGT CAGGTTTTTG TGCTTTAATT  
24541 TAATCAAGAC AATTGTTAAG GATAGCTGTG ACACTATTAT GTGTCCTTTT AATTTGATCT  
24601 ATCAATTCTT TAGAACAAAGT AATTTTTTTA AATTTAGGAA TTTTAGTCTA AAGGATTTAT

FIG. 4<sub>CONT'D</sub>



```

24661 CTTTTGGCCA TTGACAATTA GAATTTTAA TGGGGTATTT AATTCCAATA GCAACTTAAT
24721 CCAAAGTTTT CTTTATGTCA AAGAAAACAG AAGCCCAGGA GGGATGAGAC CTTGTAAGAC
24781 AAAACTCCCC TAGGAGCTTG GAATGTTTGA AAATACATGT GTTGGGCTCC CAATCTTTTC
24841 ATACTGGCTG TGATGTTACC TGAAAAATCA CATCCTTTGG ATGGTGGAGA CCAAGCGGGA
24901 ATATCCCCAT CTAGTCACGT CATGCTCTCA AGGACATGAG ACAAGAGGGA AACCTCTCAC
24961 CCTGTTTTTA TTTAGGGGAC TGGCAGCAAA GTTTGTCATA ACAGAAGTCA GCATAACCAG
25021 AACCACGAAA CTGACCAGTT TGCAGGGCCA GTTCAAACAG TGGGTTGCAG GCCTGTTCTA
25081 CCCTAGGGTA CCCCTCCTTA TGACAGAACA CCAAAAGACA AGACAAAAC GAAGGAAAAC
25141 GGCAACAACA AAAAAGCTAT TTCTGAAAGG AAAATGGCAA CAACAACAAC AAAAGCTATT
25201 TCTGAAGGGA ATGGGGTCAA ACTATGAATA CTTATACCAC AAAGTACTAA AAAATATATC
25261 AGACTCACTA TACCAAGGTT AGTCACACAC AAAACCTGTT CTCTCATTA TCTTACATTT
25321 GGAAAGGAAA AGGGAAACAA TGATTTTTAC TGTCCACTCA TCCAGAGTCC ACAGAGAGAG
25381 GAAACTGGA AACTGGGAG TCTGGCAGGA AATTCTCACT CCTCTGCTGG CTTGCCAGGT
25441 TCCTGTATTT CTTTCTCTGT GGCTTCCAGA AAAGCACAAT AGCTTTGGTG GTCTTATTTG
25501 TGATGCCAAA CTGTGGTCTT GGCCCCCTAA AGTTTCAGTG AAAATCACTG ACATGAAGCA
25561 GATTAATAGG GAAAAAGGCA TACAAATTTA TTAAATACGA ATGGGAGCCT TTAGAATGAA
25621 GCCTTGAAGC TATAGGGGAA ATTGTCTATT TTTATGTTTA GGTTTAACAA AGTATGGACA
25681 GCTGTGTAGA AATATGACTG GACAGAAAGG GCACGATCTA ATGTTAACAG ACTGAGTGGG
25741 GAAACCCAGC AAGGCCTGTC TGTGAGATT CCTCCTAGCC TCTCTCATTC CTTCTTCTG
25801 GTGTGGGGCA GGACCTCTC TGGATGGAG GTTTTATGAC CTAAGTCAAA TAACGTAGGT
25861 CAGATTTTTT TTTTTTTTTT TTTTTTTTTT GAGCTGGAGT CTCTCTGTCA ACAGGCTGGA
25921 GTGCAGTGGC GTGACCTTGG CTCCTGAAA CCTCCGCCCC CTGGGTCAA GCCATTCTCC
25981 TGCCTTAGCC TCCTGAGTAG CTGGGATTAC AGGGGTGTGC CACCACGCC AGCTAATTTT
26041 TGTATTTTTA GTACAGACAG GGTTCACCT TGTGGTCAG GCTGGTCTCA AATTCCTGAC
26101 CTTGTGATCC ACCTGCCTCG GCCTCCCAA GTGCTAGGAT TACAGGCGTG AGCCACTGTG
26161 CCCGGCCTTT TTTTTTTTTT TTTTTTTTTT GGAAGTTGTA TTTTGGGCTT TTTAACTAGC
26221 TTGTTTTTTT ATTAGATTAT TGCCCTTAGG GTGGAGCCCT TTAATAAAAA GGGGGAAGAA
26281 AACATAGGTT TTAGGGCCTC ATATTTAAAT GGGTAAAGCA GGCATAGCTG GAAGGCAGAA
26341 TACAGAACCC CCCTAATCAA GGATCTCATT TTTATATTGA ATCCTAGGCC CCCCAAAAGA
26401 GGGAAATGTC ATGGGACGAG ATGTGTGGCA TTTTATCGA GTGCCCCACT GTAAAGATGC
26461 TCCCCAAGG CTGGCAGGCA GCCCAGTGCC GATTAGCCCA CTCTGTGCTT AGTCTTTTTT
26521 TTTTTTTTTT TTTTGAGGTG GAGTCTTGCT CTGTTGCCCA GGCTGGAGTG CAATGGCGTG
26581 ATCTCGGCTC AATGCAATCT CTGTCTCGTG GGTTCAGCG ATTCTCCTGC CTCAGCCTCC
26641 CAAGTAGCTG AGATTACAGG CACCAGCCAC TATGCTCAGC TAATTTTTTG TATTTTTTAGT
26701 AGAGATGGGG TTTCAACATG TTGGCCAGGC TGGTCTCGAA CTTCTGACCC CAAGTGATCC
26761 GCGCGCCTCG GCCTCCCAA GTGCTGGGAT TACAGGCGTG AGCCACCATG CCTGGCGTGC
26821 TTAGCCTATT TTTAATGGGA GTTTCATCCT CAATGGTGAG TGCTTTCATT GTCTTTAGGT
26881 GCCCCAGACC ATGTTTTTAA AAATTTAAAT GCACGAAGAA ATAAGTAGCC CTGTATAGTA
26941 GTAATACTTT GTTGTGAATA ACTGTCATAA GTCATCTCTA AAAGTGTATT TTTTATCTAG

```

FIG. 4<sub>CONT'D</sub>

14 / 20

```

27001 TTATTATATA TGA CTAGCTA TATGTCTAGT TTTTAAATA ATACAAAGTA ATTTATTTT
27061 GGCATCCTCA AAAACCAAAG AGATTAGGTA ATGTAGTGTA GAAGAGAGCA GAGCTTTAGA
27121 CCTGAGAAGA ATCTGCCCAT GACTCGTGAA ACTCCACAAC GAAAGTAGGA GACCCCAAAA
27181 AAGGGGTGAG TGTCATCTTT TCTGAATTTT TTTTTTTTTT TAGATGGAGT CTTGCTCTGC
27241 CACCAGGCTG GAGTGCAGTG GTGCAATCTC GGCTCAGCCT CCCGAGTAGC TAGGATTACA
27301 GGCACGCGCC ACCATGACCA GCTAATTTTT GTATTTTTTAG TAGAGACAGC GTTTCACCAT
27361 GTTGGCCAGG ATGGTCTCGG TCTCTTGACC TCGTGATCCG CCCGCCCTCGG CCTCCCAAAG
27421 TGCTGGGATT ACAAGCGTGA GCCACTGCAC TCGGCCCGGC AGATAATTTT TTTGGCCAGT
27481 TTTTACATAG AGTAATTTTA GGTTTTATGG CTGGCTTTGG GGCAAAGGGG TTCTGGTTTT
27541 TATAGCTGGT CTTGGGGGAG AATGGAACCG AGTGACAAGA GGACAAGAGA GGGTCAGAGA
27601 AAAACTTCTG CTTCTGAGGC GGCTATTGAG GCCTTCATTT TGGAGTATTG TCCTCTAAGC
27661 CCCAGCAGTG TCAAACGTGA CACAAACCAT ACACAGCAGC CAGCTCGGGT GCTGTTAGGA
27721 AATGGTCTCA CTGCTGGGTC TGTGGGGTAT GTGTGTGTCT GGGTGTGTGG CTACTGTCTG
27781 CATCCTCCTC CCCCTACAG CCTCCCCGCC TCCCCTCCAG CCACCCTGGG ATTGGTGA CT
27841 CTCAGCCCCT CCCCTCAGCT CCCCTAGACC CTCCCAGAGC CTTTATCAGG GAGCTGGGAC
27901 TGAGTGACTG GAGCCTTCCT AGATCCCCCT CACTCGGTTT CTCTCTTTC AGGAGCACCG
27961 GCAGCACCAG TGTGTGAGGG GAGCAGGCAG CGGTCCTAGC CAGTTCCTTG ATCCTGCCAG
28021 ACCACCCAGC CCCCAGCACA GAGCTGCTCC ACAGGTAGGC AAGTGGGAGA ATGCTGGATG
28081 GACCAGAGCT GGCACCAGGG GACAGGAGCC AGCGTCAGGA GGAATAAAG CAGATGGCAG
28141 CCTCTGATAG GGGAGCAGGG GACTGGGAAG GTGAGCACAA AGCACCTGTA GGGCCGAGAG
28201 CTGGTTGGTG TTTGGAGCCT GTGGCTACAG ACTCATCTT TCATACCAGA AAGTTTTTGC
28261 CTAAGTCTTG GGATTATCTA GTACTGGAAA ATAGCATCCA GGATCCCTCC TCCAGCTGAC
28321 TGAGGAAACA GACCAGTCCA TGTCTACAA ATCTATCATC TTTCTTGGGA GCTAGAGTCC
28381 TCCTGGCACC ACTATAGCAT TGCACATCTC CTGGGGAGAT ATCTGATGGG GTAGCAGGGA
28441 AACTAAGCCC AAGGGCTGTA CCCCCTTCTC AGAAATACTT TCCACCCTCT CTCCAGACCA
28501 GGGCTTGGAC AGTGGAGTTG GGGGCTGGGG AAGCAGGGTC AAGCCAAGCT GCTGGTAATG
28561 AATGTCTCTT GTGTCTTCAC CCATGCTGTA TCTTCCTCTT CTCTCCTTTA CCTGAGTCCT
28621 GTCCCTTTGC TCTCCAGGC ACCATGAGGA TCATGCTGCT ATTCACAGCC ATCCTGGCCT
28681 TCAGCCTAGC TCAGAGCTTT GGGGCTGTCT GTAAGGAGCC ACAGGAGGAG GTGGTTCTTG
28741 GCGGGGGCCG CAGCAAGGTA AGTCTCCCTT GGCAGAGTAC TGGGGACATC ACGGGAACCT
28801 GGGACTCTGC CTGTCTGGAC AGCTGTAGTG AGGAAACTGG GGTGGGGGGG TTGTCCGTCA
28861 GAGGGCATT TGCCTCCCTT TGGATTTCTT TGTTTCTCTG GTCCTTTCAT GTTCCCACTG
28921 TCTCCAGGTG TGTTTGTGTC TCTGTATCTC TGCATGTCTT TGACACCTTG TACATAAAAG
28981 GTGCCCTACA AATATGTTGT TTGGTGGGTT GATTGATGGG AGACTTGGTG ATTGGATGGT
29041 ACTGTGAGGG GTGAGCTAGG GTGGTCTAAG GCTCTCTATA GTCTACCTCA GGTCCCTTTG
29101 CAAGGGACAG ATCTCTTCTA TTTCTGGGAT GGTATGAAAC AGTCAGAATT TCTTTCCCAA
29161 ATGGTTATTT GTGTGCTATT TTACCTATCA GTTATGTGTA TTGTTTTATT TTCAAATGC
29221 AAATAAATTC CTTATCTTT TGCTCATCCA CCCCAGTAA CCTCAGGTGC TTCTAAGATC

```

FIG. 4<sub>CONT'D</sub>

15 / 20

```

29281 CCAACCCCTT CCTTCTTCTC TTTCTCCCTC TGCCCGCCTC TATCCTCTGC TTAGTCAGGA
29341 TAGGAAAACA ACAACAGCAA AAAAACCAGA TTGAGCCTCG ATTTCCACAG TTCCTTTACG
29401 AAAAAGAATA GGAATTGTCA GGGTAGGGGT ACAGGGGGAG GATAGGGAGG AAGTCTTTTC
29461 AAGGTTTTGA AATGACAGCA ATTACATCGG TACAAATGCT TTTAAGATGA TTGCGGGTGG
29521 GACTTATTAC AAATTCAATG TGTGAAGTTT AACTGCCTCT TCAGCTCAA TCTGTTTCAGC
29581 ATCTCATTAT AGGAGGTGGG CAGAGTATTC AACAATTTGG GAAAAGTGGC TGCCTGAACA
29641 CCACATGCTG GGCCAAGGGA GTTATCACCA GGGCAGCCTT GCAGGTGGCA GCAGTTGTGC
29701 CATATCCAAA AGGCCAGAAC CGTTAAAAAA AAAAACACCC AGGGGAGTGC CAAGTATGGG
29761 CTGGACACCG TTTGGAGCCA CAAAGTTCCA GCCCAGGATA GTTAGAGTAT CTGAGTTCTT
29821 CTGAGACAAA CTTGTTTCAA GACCTTGGCC AATGAGATGT CCCCTCTGCC CCTCTTGGTC
29881 AATGAATGAG AGGGATTGCC ATCCTACCCC TTCTCCTTGA GAGTCTGTGA GGATGAGGGA
29941 AATTGGGGCA GGAAGAGGGT AGTACATAGG TGTGCCTAGG CAACTGGGTT GGTATGTGTG
30001 GGGGTGTGTT CTGTGTAAAT GCACCTCTGT GTGTGCACAA CAGCCGAAGG ATGCCTGGGT
30061 TCTGGAAAGA GAGGCGCTGC TGAGACTTGA GATTTGAGAT GAAAATCTCC AGCCATGATC
30121 ATTGTTATTG TCTCTCTGCA GCTGCAATTA ACTGGCTGTG TGGTGTGTGC CCACCACCTT
30181 GCTGTACGCA AGTTGCTAAA AAAAAAAAAA AAATCACAGG GACAATCAAG AGCCCGTGCT
30241 GGGCAACAGC TCTAGAACTT GGGATTCACT TGTGGAGAGA AGAAGACGTG CCTTCTGAGC
30301 ATGTTGCCTT CCTGGAATTC TAGACCTAGG GCCAAAAGGG AGAGGGAGAG AAAACTAGAG
30361 GCGGAAAGCC ATGGAGAATA GAGAAAGAGG TGGTGGAAAA CAGGGAGAGA AACATCCATG
30421 GACATCGTGC AGAGTGGGGG AATCACAGGT GCAGATGTGT GCCTCCAATC TCACCATGCA
30481 TGTGAATCAC CTGGGGGGCT GCTTAAATATG CAGATTCTGT CTCAGGAGGT CTGGGGTAGG
30541 AACAAGAGTC TGCATTTCTA ACAGGCTCTG TGTAGTGCTG GTGTTGCTGT TGGTCCACAG
30601 GTCACCTCTG GAGCACCTAC TTCTCGTCCA GTGTGAACCA GAGGAACTC TGAAAGAAAT
30661 AGGGTGTCGG ATTCAGGATG GGCTCAGGAA GAGGCTGTTT CTTGTGGGAA AAGGATGAGT
30721 GGATCCGGGT GGGAGCCTCC TGCCTCACCC CTCTTTGTTT CTTCCCTAGA GGGATCCAGA
30781 TCTCTACCAG CTGCTCCAGA GACTCTTCAA AAGCCACTCA TCTCTGGAGG GATTGCTCAA
30841 AGCCCTGAGC CAGGCTAGCA CAGGTAGGAG GCGGCCCTAG GGGAGAGGGG AATGAGGGGC
30901 AGGATTCTGA AGATAAGAGG CCTGGGAGAT CCTTTCAGAT GGGAGAGAGA TGGGGGATAG
30961 CTTAGTGAAT CCGTGAGGGT TGTGATCTGA ACCCCGCTCT CATCACTTTC CAACTTCACT
31021 CCCCATTTAG ACATCTGTTC TTGGTTTCAC AGATCCTAAG GAATCAACAT CTCCCGAGAA
31081 ACGTAAGTAC CCTCTTCTCC CTCCCTATCT CTTGCCACTT GCCCAGAGCT CTGTGGGGCA
31141 TTGGGCCCAG GGGCCATTTT GTCCAGCCCC TTCTCACCTG GTACAAACAA TATGCCAGCT
31201 CCCACTGCTC AGCCAACCTT TCCTGAAAGG GAGAGGCCAT CCAGAACTAG GAGGAAGCTG
31261 GTGTGAGGGG CATGGTGGGC TCTCCCTCTG CTGGCTGGTC CTTGGAAAAC AAGGGGATCT
31321 CTTCTGTGGC CTGAAAATTC CAAATCAGGC ACCTGCTAGA GCAGAAAATT CTTGAAATGT
31381 GGAGGAAGGA AAGGTGAGCA GAGAGAGTGG GTTTAGGGGA GGCCTTGCT AACTGTGAGG
31441 AGTCATGCTT TGACAAGAAA AAGGAACAGA GACCAGAAAC CCAGTCTCAG AAGTGTTGAC
31501 CCATGTCTGG GGAGATGCTT CACTTTCTCA TCATCACTGC TGACAATGTT GGCCCTTTTC
31561 TGCAGGTGAC ATGCATGACT TCTTTGTGGG ACTTATGGGC AAGAGGAGCG TCCAGCCAGG

```

FIG. 4<sub>CONT'D</sub>

```

31621 TAGGAGTGTG TGGAGGTACA GTGGAAGGGC TTAGGGTACT GGCAGAGTAT GACAGAAGTC
31681 ACGTGCCTCA TATTTGTCAC CAGAGGGAAA GACAGGACCT TTCTTACCTT CAGTGAGGGT
31741 TCCTCGGCCC CTTTCATCCCA ATCAGCTTGG ATCCACAGGA AAGTCTTCCC TGGGAACAGA
31801 GGAGCAGAGA CCTTTATAAG GTAGTCTGTG TGCAGCTGGG AGGAAGGATA GGGAGACTCT
31861 GCTTCCACCC CAGTCTCCCA ACTCTGTCTT TGAACACTGC CCGTCATAGC CAGCCCTTTG
31921 CTGTTGGATC AGGGTGTAGT TCACATTCAG AAAGATCCCT CTTACTTACA CTGTTGCTT
31981 TACCCTAGAC TCTCCTACGG ATGTGAATCA AGAGAACGTC CCCAGCTTTG GCATCCTCAA
32041 GTATCCCCCG AGAGCAGAAT AGGGTAAGGA TTGTTTATTA GAGAGGGGAG AGGGGACTGG
32101 GGAGGGGGCT GTGGGGGTTG CCAGCTGTGC ATTTCTCTCC ATGCTACAGG TATTAAAGCT
32161 CATAGATTTG CCCTGAAATA CACTGCCAAT GCCCAGCACA CTGTCGGCCA AACACAAAGA
32221 CACTTAGAGG CACGTGTGTT TGTACACATC CCCCCTCTTT CATCTCTTTC CTCTGGATCA
32281 TGGACGGCAG CTGACTATTG AGCAGGAGTG AGTGTGGGA GATGAGGAGA GAGGGGCTTC
32341 CCGATGGGCA ATTTCTGTTG TTTGGACTTC ATTCTTTTGT AATCTATGCA AAAAGATGGA
32401 GAAATTATTA TCTGATAATT ACAAATACCA CAACCAATTC ACAGGCAAGC ATTTGCCTCC
32461 CAGGCAGGCT GAGCCTTTCA AATCACTCAG AATCCTGGGT TACGGGGCCC AGAAGGTAGT
32521 CATACACAAG GATGATTCAG GAAGAAATGC AAGGAACTCT GAAATCTAAT GGGGATTAGC
32581 AGGAAACCAT ATCTGAATCT CTCTTTAGCA TAATGAATAA GAACAATGGC CTGAATGTGA
32641 ATCCTGGATC TGCCACTCTA TCTGTATCTT TTTGGCCAAG GTACATATCC TCCTGTGCTT
32701 CAGTTTCCTC ATCTGAAAAA TGAAAGTGAT AATAGTATCT CACAGGGTTG TGGTTTTGAG
32761 GATTGAGTAT AGGTAAAGTG TTCAGAACAG TGCCGGGTGC ACAGTGCTGT GTGCCAATTT
32821 TATGATAATT GTCCCAGTTT GGGAGGTATG GGGGATGTCC TAATGTTTCC CCTGACTGGC
32881 TCTGTCTGGA CCCCAGGCCT GAGTGGGCTG ACAAATTCCT CACTTGGTAT GCGAGTGTA
32941 GAGTCCCCCA GGAAGTGTC TAGTCAAAAC ACGAACCTTC CGCCTTGACA CTGTCTTCCC
33001 ACACACAGCA AGAGCAGCTC CACCAATGGC TTTCTTTTCA CTAGCTTCCA AAGAATTGGG
33061 GTGGAGGGAG TGAAAAGGAG AGGGAGAGAG ATTGGGAAGG CTCGTAATCA TGGAGAGCCT
33121 CCTGCTTTTC TCTCTGTGTC CCTGTTACCC ATACTCACTG GTCTCAAGGT GGCACGCCCA
33181 AGACCCAAGG AGCTGGTGCT TGATGATGCT GCCTGTGCAT GAATTCCTGG GACCAGAGAC
33241 TGAGTCTGGC CCCCCATTTA GTGTTGGGTG AGAGGGCACA AAGAGCTATA ATAAGTGTA
33301 CTTGCTGATT ACATGGTAGT TACTGTATCA TTTTGCTCTC ATTAGATGGT TATTTAGTCT
33361 CTGCCGACGG CCAGATAAAT ATACGAGCAG CTATATCTGG ATGACATACT CTGCTCCAGC
33421 GTTATGCACT GGCCATAAAG ATAATTACAG TGCAATTTTG CTATAGTATT TTATACAAAT
33481 GGCAAAAACA AGTGCATTGT GGAAATCTAC TTTTAATGCT TGTTTGTGCA TCCAGGCTCT
33541 TTCAGAGGGA CCCATAATTG CAGCTTTCAT AATCTTACCA TTGAGGGAGC ATTTCCCAACC
33601 TGTTAGGTGT CAGGCAGAAT AGGACATAAG GTTTCTGGGA GCTGGCATTT AAAGATTAGA
33661 TGAGATGGAT CAACACAGAT CATTGTGTCA TCTGATTTCA TTCATGTGAA ACTGTAAGTA
33721 ATCCCTGGGC CTGTGCTTCC TCTGGGAGGT TTCTGGGAAG AGGAGGAACT GGATAAGGCA
33781 GGGGGAGCAT TCATAGTAGG GCACCTTGGG CAGGGCTGTG TGTGTGCTG GCTCATGGTG
33841 GTGCTAGGAT GGCATGAACT TGGTTCCTAC ATCTTTGGTC CACATGGGCC CCACTGGCCA

```

FIG. 4<sub>CONT'D</sub>

17 / 20

33901	TGCACACAGG	TGTGTAGAGT	AATGTAAATA	TGGCAGCTGG	GAAGGTGCAA	GTACCTGCGG
33961	CTAGGAGAGT	TCCATCCTCA	GGCCCAAAGC	CTGGAGGGCA	GGCTGAGGGT	CAAGACTTGT
34021	TCTTTCTCT	CTCACAGACG	CCTCTCCCCT	TCTCTCCTGC	TGCCACAGCA	GGTTTTCAGT
34081	GGGACTTTTT	TACAGGATAT	AAGATGTGAT	TTCAGTGTTT	TTTTTTGTTT	TGTTTTGTTT
34141	TTTGTCTCA	GTACTCCACT	TCCGGACTCC	TGGACTGCAT	TAGGAAGACC	TCTTTCCCTG
34201	TCCCAATCCC	CAGGTGCGCA	CGCTCCTGTT	ACCCTTTCTC	TTCCCTGTTC	TTGTAACATT
34261	CTTGTGCTTT	GACTCCTTCT	CCATCTTTTC	TACCTGACCC	TGGTGTGGAA	ACTGCATAGT
34321	GAATATCCCC	AACCCCAATG	GGCATTGACT	GTAGAATACC	CTAGAGTTCC	TGTAGTGTCC
34381	TACATTAAAA	ATATAATGTC	TCTCTCTATT	CCTCAACAAT	AAAGGATTTT	TGCATATGAA
34441	TGATGTGGTG	TGTGTGTTTA	CTTGTTTGGT	TGGTGGGTTT	TTCTGTTCTT	TGACTCCTCC
34501	AGCTACATGG	TAAATACACA	CATACTTATG	ATACACACAC	TTCATATTTA	AATGTTAAATA
34561	ACTTTACATA	TCTTTTTGTA	TATATCTATT	TCCTGAACAG	TGCCTTACAC	AGTGCTTTGC
34621	ACGATGAGTA	TCAGATTTAT	TTAGTGATTA	AAATAAATAC	ACGAATTTGG	AAGATGGTTT
34681	CTAACACACA	AAGATTTTTA	CAGACCAGTT	TTAGATAAAG	AAAAAACAGG	CCGGGCCCCG
34741	TGGCTCACGC	CTGTAATCCC	AGCACTTTGG	GAGGCCGAGG	CGGGTGGATC	ACGAGGTCAG
34801	GAGGTCGAGA	CCAGCCTGAC	CAACATGGTG	AAACCCCTTC	TCTACTAAAA	ATACAAAAAT
34861	TAGCCAGGCA	TGGTGGCGCA	TGCCTGTAAT	TCCAGCTACT	TGGGAGGCTG	AGGCAGGAGA
34921	ATCGTTTGAA	CCCAGGAGGC	AGGGGTTGCA	GTGAGCCGAG	ATCACGCCAC	TGCACTCCAG
34981	CCTGGGCAAC	AAGAGCAAAA	AACCTCCGCT	CAAAACAAAA	CAAACAAACA	AAAAAAACAA
35041	TAAAAAAGA	AAAAGAAAAA	GAAAAAAAT	ATTGAGAATG	ACTTGATTTA	CTAGGATGGG
35101	TCTGGGAGAT	ATTCAATCCCT	GAATCTGACC	CTACTTAATT	AGAGAAGGAG	GTGGGGATCA
35161	AGGCTGTCCG	GAGACCCAGC	CACAGAGGAG	GACAAATCTA	TGACCCATATA	CAATTTTTTT
35221	GTCTCCAAAT	GCTGAGCCTG	GGTTCTGTGA	CAGATCCTGG	GGATGAAATG	ATGACTCATA
35281	CACAGAGTTT	ACAGTTTAGC	AGGGCTGTGG	ACAAGCAAAC	AGAACTTGAT	CCAGCTAGGA
35341	TGGGATGTGG	ACAGGGAAGT	TACTACCGAG	GCCAAGAAAG	AGAGGAGCAG	ATATCTTCAC
35401	CGTTAACTGG	CTGCCTTAGT	TATTATAAAG	GGAAAACATT	TATCTCCAC	TCCTCTCTAA
35461	AGTGCCTGTT	ACCAGCTCCT	GCAGCTCTGA	CTTAACAGTC	CCCAGAAATG	GTAAGGCACT
35521	TACATGTGGT	ATGCATGGGT	ATGGATGTCT	TTTACTAATC	TATGATGTCA	ACTATCACCC
35581	GCCATCCTAA	GGGGGGTTCT	GTACCCTAAT	GGAACAGCCA	GTGAAATCCT	CAGGCTCCTT
35641	ATCTTAGCGT	GGTACAGGGG	CCTTTGTTAT	GCCCCTGAAT	TGCACTGATA	AAACATCAAC
35701	ACATAGATTT	CCCAAGGCAG	TGTAAGGACA	GGGCCACAGA	GCCAGAGGCC	ACTTCCTGCA
35761	GTCCTTTTCA	TCTAGTGAAA	ATTCTATCTT	CCTACAGCCT	GACTTGGGGC	CACCTTTGGAA
35821	TGACAGCTGT	ATAGTGGGGG	GCGGGGAAAG	GAGGGAATAC	TCACCCTAGT	ATTACTTATG
35881	TCAGCTTTAT	AGCCAGAGGT	CAAAGAATGC	CCCCACCCCA	GAGCCTAGAC	CCTTTTTTCCA
35941	GTGAGTCATC	TCTTTGACTT	TTCAAAATTA	TCTATCTATA	GGGCTTAAAA	CTGGGGACAC
36001	TTTTGCAGAG	TCTAGGGGCT	TTCTCTGGGT	CATGAAAGCT	ACAAGAGTTG	GTTCTGCTCA
36061	GACTTGGTGG	GAGTTAGGCT	TATAGGCTGA	GATGAGACAA	TTGCTTTGCA	AGTAGGAACA
36121	TTAAGTGCAG	AAAGATTGCT	CTCTAGTGGG	ACTGACAAAA	ATTGCAGTAC	TGGGGACTCC
36181	AGAAAAAAT	GAAGACAAAT	GTTAAGTTAG	ATTCCTGTGT	TTGTACTTGA	AGAATGTGTG

FIG. 4<sub>CONT'D</sub>

18 / 20


36241 AAGGGATCCT GACCCTCCCT TTCCTGTTGT AAAACAGTTG ATGCCTAAAG AGATCTGGTC  
36301 CACAAGACCT TGAATAAAT CCTGGCCCTT TCTTCTCCAT TTAACCTTGT ATATGTTTGT  
36361 TATTGTGACT ATATGGTGAT TTAACCTTAAA AAGACTTCAG TATAAGTGGT ATATACTTTC  
36421 ACCTGCGTCT TTTGGATGAT TTGTTTTCAT GTGAAGTTTA TTGGGGTCAA CCCTCCAGAG  
36481 ATGGCTGGGG CAGTTGGTTA GAAAGACTGT ATAGGCCAG GCCCTTGCAA GCCCAGCAGC  
36541 CCTCTGTCTC CAGAGTCATG CTGGAGGTCT GGACCTGCTG GCTGTGTGAT ATTCCACTTT  
36601 AGGGAGACTC AGTCACCTTG CACAACCTGTG AGAGCTGGGC CTGCCACTGA AACATTGTGT  
36661 CAACCTCTAA GTGACCCTTT CACTAGATGG TAAAGTGAGA TGCCTCATCC CCAAACCTATA  
36721 AGAACAGTTC TATGGCTGTT TTTGTATCTC CTGGCTAACA AATGTTACAT GTTTGGCAGC  
36781 ATTTGGTATA GTGCTTGCTT TCAGTATAGT CTGCCACCAG TTAATGAGGT TGTGGAAAGG  
36841 AGGACACACA ATCTCCCAA TTCATCAAGA GAATGGACAA TTGCTGAATG GCCAACTGG  
36901 CTTAGATCTG TTGGCAACAT TCAGTGTGTC CCTTCCTTTC CACTTATCCA TCAAGGAATT  
36961 ACTGAATCCT ACCATGCGCC TGTCTGGGA GTTTGTCCTT GGCTGCAAGC TATTTTCAGG  
37021 CAGTGACTGG GATGGGATGG GAGAGAGGAT GAAACTGAAG GGTCTTGGAG CCTAAGAGCT  
37081 TCCTCTGTAC TGAGGGAGGG AGGGCGACAT GACGAAGACT TCTAATGTCT TTGGTGGTGG  
37141 TGGGTGGGGC AGGCAGTGTA GGTGGTTTTT GTTTGATGAC AATTCCTGGG CAGAAGCATT  
37201 TGAAAAGATG ATTTGGGAGA AGGGTGGGGA GGAAGAGTGA TCGAGTTCTA CACAGAGTTG  
37261 GGGAGGGCAG GCTTCAGGAA GCAGGCCTGG GGTGCCAAAG TACAGTGAGA TCCGGTGACT  
37321 TTCTTCATTT GGCCACCTAG ATGGAAGGAG GGACAGCAGT GGATTATCAG AAGGGTCCAG  
37381 TAGTAGCGGT CTAGCCCTCA AGTGCTCCTT CATTCATTCA AGCAGGCTTA ATGTATTAAG  
37441 CACTTATTGT GCCAGGAAGT GTGGTAAGGG TCAGTGTGGA CCTGCGGCCG TGTGCAAAGC  
37501 CACAGATCCC TGCCTTCAGG AAGCCCACAG CCTAGTGGAG GAGATATATA GTAATCAAAC  
37561 AATCTTACAA CATTTTGTAA AATGCCATA GTAGATGTTT TGAGGAGAAG CTTTTGGAAC  
37621 TGTGAGCGTA GAACAGGGGA GGTGAAGAGA GTTTGGATAG G

FIG. 4<sub>CONT'D</sub>

19 / 20


## FIG. 5

QUANTITATIVE PCR OF THE COMPLETE HUMAN NEUROKININ B PRECURSOR



## FIG. 6

HIGH PRESSURE LIQUID CHROMATOGRAPHY (HPLC) OF NEUROKININ B



20 / 20

FIG. 7

CARDIOVASCULAR EFFECTS OF NKB IN CONSCIOUS RATS




FIG. 8

LOCALISATION OF NEUROKININ B mRNA EXPRESSION  
IN VERTICAL SECTIONS OF THE PLACENTA